


# TOXIC AIR: THE PRICE OF FOSSIL FUELS

June 2020

GREENPEACE  
غرينبيس

# CONTENTS



## Authors:

Aidan Farrow, Kathryn A Miller and Lauri Myllyvirta

## Contributors:

Erin Newport and Minwoo Son

## Acknowledgments:

CREA - Centre for Research on Energy and clean Air

## Design:

Jihane Abou Khater

## Published by:

The original version was published in February 2020 by Greenpeace Southeast Asia.

This version is edited and released by Greenpeace MENA.

For more informations contact: [info.arabic@greenpeace.org](mailto:info.arabic@greenpeace.org)

## Suggested citation:

Farrow, A., Miller, K.A. & Myllyvirta, L. Toxic air: The price of fossil fuels. Greenpeace Middle East and North Africa 35pp. June 2020

<b>Executive summary</b>	<b>01</b>
<b>1.0 Introduction</b>	<b>02</b>
1.1 Air pollution: a brief overview	02
1.1.1 What are the key air pollutants?	02
1.1.2 What are the sources of air pollution?	03
Natural	03
Anthropogenic	03
Primary and secondary pollutants	03
1.2 Air pollution from burning fossil fuels	03
1.3 Air pollution, health and cost	04
1.4 Air pollution and climate	04
<b>2.0 The economic cost of air pollution from fossil fuels</b>	<b>05</b>
2.1 Introduction	05
2.2 Health impacts and costs	06
2.2.1 Health	06
2.2.2 Economic cost	09
2.3 Regional examples of the cost of air pollution	10
2.3.1 Health	10
2.3.2 Economic cost	14
<b>3.0 What can be done about anthropogenic air pollution?</b>	<b>17</b>
3.1 Case study one: Switch to sustainable transport	17
3.2 Case study two: Generate electricity from renewables, not fossil fuels	19
<b>4.0 Conclusions</b>	<b>21</b>
4.1 Costs	21
4.2 Transport	21
4.3 Energy	21
<b>Glossary</b>	<b>22</b>
<b>Appendix 1: Methodology</b>	<b>23</b>
Deaths of small children from lower respiratory infections	23
Diabetes, asthma and other chronic respiratory diseases, and disabilities caused by stroke	23
Preterm Birth	24
Work Absence	24
<b>Appendix 2: Summary of cost and mortality data</b>	<b>25</b>



## EXECUTIVE SUMMARY

**This report reveals the cost of air pollution from fossil fuels and highlights solutions that can protect our health and benefit our communities. Air pollution generated by burning fossil fuels is attributed to approximately 4.5 million premature deaths worldwide every year, the report shows. Air pollution increases the incidence of chronic and acute illnesses and contributes to millions of hospital visits and billions of work absences due to illness each year. It also damages our economies and the environment.**

For the first time, Greenpeace Southeast Asia and the Center for Research on Energy and Clean Air (CREA) have quantified the global cost of air pollution from fossil fuels, finding that it has reached an estimated US\$8 billion per day, or 3.3% of the world's GDP. While coal, oil and vehicle companies continue to push outdated technologies, our health and our communities are paying the price.

The economic cost of air pollution reflects pollution concentrations, population size and the availability and cost of healthcare. We found that the China Mainland, the United States and India bear the highest costs from fossil fuel air pollution worldwide, at an estimated US\$900 billion, US\$600 billion and US\$150 billion per year, respectively. For the Middle East and North Africa region (MENA), Egypt, Saudi Arabia and the United Arab Emirates bear the highest costs from fossil fuel air pollution at an estimated US\$6.9 billion, US\$6 billion and US\$5.9 billion per year, respectively.

We estimate that exposure to PM<sub>2.5</sub> and ozone from fossil fuels is responsible for 7.7 million asthma-related trips to the emergency room each year globally and 550,000 in the MENA region alone. Exposure to fossil fuel generated fine particulate matter (PM<sub>2.5</sub>) alone is attributed to an estimated 1.8 billion days of sick leave annually, 40 million of those are in the MENA region.

In the MENA countries, the number of premature deaths annually is estimated at 65,000 as a result of air pollution caused by fossil fuels. Egypt, Lebanon and Syria recorded the highest estimated premature death rates due to air pollution from fossil fuel consumption in the MENA region.

Air pollution is a major health threat to children, particularly in low income countries. Worldwide an estimated 40,000 children die before their fifth birthday because of exposure to PM<sub>2.5</sub> pollution from fossil fuels. In the MENA region, the estimated number is 865 children. We found that air pollution from fossil fuel-related PM<sub>2.5</sub> is attributed to an estimated 2 million preterm births each year.

Yet while toxic air pollution is a global threat, the solutions are increasingly available and affordable. Moreover, many solutions to fossil fuel air pollution are also the solutions to climate change. Clean transport and renewable energy not only bring significant reductions in toxic pollutants such as PM<sub>2.5</sub>, NO<sub>x</sub> and ozone, but also help to keep climate change-causing greenhouse gases out of the atmosphere.

A phaseout of existing coal, oil and gas infrastructure is not only essential to avoid the worst impacts of global climate change, but it brings major health benefits due to the associated reduction in air pollution. Research shows that the closure of coal-fired power plants can yield health benefits that exceed the value of electricity generated<sup>1</sup>. According to a study published in the Proceedings of the National Academy of Sciences, an expanded fossil fuel phaseout and investment in clean energy sources could reduce premature deaths related to air pollution worldwide by up to nearly two thirds<sup>2</sup>.

In addition, a transition to affordable and carbon neutral transport is critical to ensuring healthy cities. Effective public transport systems and good walking and cycling infrastructure enable mobility, reduce air pollution and greenhouse gas emissions, and correlate with a decrease in rates of cardiovascular disease, cancer, obesity, diabetes, mental illness, and respiratory disease<sup>3</sup>.

One of the most important ways that governments can catalyze sustainable transport is to set a phase-out date for diesel, gas, and petrol cars, and to introduce comprehensible and affordable public transport, with safe walking and cycling infrastructure. We need to move away from private cars as the primary mode of transport, and initiatives like car-free days allow us to imagine what our cities would look like without traffic and pollution.

The transition to renewable energy is essential both to prevent catastrophic climate change and to protect our health. While fossil fuel companies continue to market outmoded technologies, our communities pay the price. A just transition to renewable energy is possible, but we can't afford to delay any longer.



# 1.0 INTRODUCTION

This report, 'Toxic air: The price of fossil fuels', assesses the impacts on global health and the economic cost of air pollution from the continued burning of fossil fuels such as coal, oil and gas. Using data published in 2019 – including the first study to assess the contribution of fossil fuels to global air pollution and health<sup>4</sup> – the report provides a global assessment of the health impact of air pollution from fossil fuels in 2018 and a first-of-its-kind estimate of the associated economic cost. Case studies relating to transport and power generation show that reducing air pollution is feasible, achievable and cost effective.

This report uses the most recent evidence and data on pollution levels, health effects and demographics to quantify the effects of air pollution on global and regional levels. The analysis includes an estimate of the financial cost of the health burden on the global economy. Using case studies, the report discusses how the phaseout of fossil fuels will have the co-benefit of mitigating climate change and reversing some of the most pressing global health problems<sup>5</sup>.

## 1.1 Air pollution: a brief overview

The primary focus of 'Toxic air: The price of fossil fuels', is the impact of air pollution from burning fossil fuels on human health and the associated financial costs. The study is limited to the pollutants; fine particulate matter (abbreviated to PM<sub>2.5</sub>), ozone (O<sub>3</sub>) and nitrogen dioxide (or NO<sub>2</sub>), and only that pollution which is emitted by fossil fuel combustion.

### 1.1.1 What are the key air pollutants?

This report only considers the impact of fossil fuel-related air pollution and only those pollutants for which there are well understood relationships between changes in pollutant concentration and health impacts. The following pollutants are included:

**•Nitrogen oxides.** When fossil fuels are burned in air, nitrogen oxides (NO and NO<sub>2</sub>, collectively referred to as NO<sub>x</sub>) are created from molecular nitrogen in the air and in the fuel that is being burned. NO<sub>x</sub> pollution, along with sulfur dioxide, which is also produced when fossil fuels are burned, reacts with water to form acid rain, snow and fog, and with other substances to form particulate matter (see section 1.1.1) and smog. The health impacts of exposure to nitrogen oxides include cardiovascular diseases, exacerbated symptoms of asthma, chronic obstructive pulmonary disorder and other respiratory diseases<sup>6,7</sup>. Acid rain is detrimental to plants and animals.

**•Ozone.** Ozone (O<sub>3</sub>) is found in the stratosphere, one of the Earth's protective atmospheric layers. Stratospheric ozone protects the Earth's surface from ultraviolet radiation from the sun, but ozone also forms at near-ground level, where it is an air pollutant that causes smog. Ground level ozone forms when NO<sub>x</sub> pollution reacts with chemicals called volatile organic compounds. Ozone pollution causes acute human health problems, including chest pain, throat irritation and inflammation of the airways. Ozone can also impair lung function and increase the symptoms of bronchitis, emphysema and asthma<sup>8</sup>. It also adversely affects vegetation and crops.


**•Particulate matter.** Particulate matter, (also known as particle pollution or PM,) is a term used to describe extremely small particles and liquid droplets in the atmosphere. In relation to exposure to particulate matter, the World Health Organisation says: "There is no evidence of a safe level of exposure or a threshold below which no adverse health effects occur"<sup>9</sup>. These particles can be a combination of different chemicals and are classed according to particle size: PM<sub>10</sub>; PM<sub>2.5</sub>; and ultrafine particles (Figure. 1).

**a) Ultrafine definition.** Ultrafine particulates have no formal definition but the general consensus is that they are any particles with an aerodynamic diameter of  $\leq 0.1 \mu\text{m}$ . Ultrafine particles are respirable, which means that they are small enough to reach the gas exchange region of the lungs<sup>10</sup>.

**b) PM<sub>2.5</sub> definition.** PM<sub>2.5</sub> refers to any particulate matter with an aerodynamic diameter that measures  $\leq 2.5 \mu\text{m}$ , including ultrafine particles. As with ultrafine particulates, PM<sub>2.5</sub> particles are respirable, which means that they are small enough to reach the gas exchange region of the lungs.

**c) PM<sub>10</sub> definition.** PM<sub>10</sub> particles are  $\leq 10 \mu\text{m}$  in diameter<sup>11</sup>. These particulates are inhalable and can lodge in the respiratory tract<sup>12</sup>.

Figure 1. A: Graphic representation on a logarithmic scale showing example sources of different sized air pollution particles (blue, red, yellow, green) and the size fractions of particulate matter to which they contribute (dark to light grey)<sup>13</sup>.



### 1.1.2 What are the sources of air pollution?

#### Natural

Particulate matter occurs naturally in the environment. Airborne desert dust, sulfates, volcanic emissions and organics released by vegetation are natural sources of PM<sup>14</sup>. Nitrogen oxides are released into the environment from natural sources such as microbial processes in soils, lightning and forest fires<sup>15</sup>. Many of these processes can be exacerbated by human-induced global warming and environmental changes.

#### Anthropogenic

Human activities that contribute significant quantities of particulate matter include road- and non-road transport, including shipping and air traffic; public energy production by fossil fuel power plants; commercial and residential combustion sources (cooking and heating); industrial activity; biomass burning (forest, shrub, grass and agricultural waste); and agriculture. In urban areas, traffic and combustion are the primary sources<sup>16</sup> of PM<sub>2.5</sub>.

Nitrogen oxides are released during any combustion reaction, particularly at high temperatures. The primary anthropogenic sources of NO<sub>x</sub> are vehicles, non-road vehicles, (for example, construction equipment,) industrial sources such as power plants, turbines, industrial boilers and cement kilns, boats and heating for buildings<sup>17</sup>.

#### Primary and secondary pollutants

Air pollutants are classed as primary if they are emitted directly from the source – a factory chimney or vehicle exhaust, for example. Secondary air pollutants are formed when a chemical reaction occurs in the atmosphere involving a primary air pollutant. Ozone is a secondary air pollutant created when oxides of nitrogen react with a group of chemicals called volatile organic compounds. Some particulate matter pollution is secondary pollution. For example, sulfur dioxide can oxidise to form sulfuric acid, which can then produce ammonium sulfate particles if it reacts with ammonia.


## 1.2 Air pollution from burning fossil fuels

Burning fossil fuels – primarily coal, oil and gas – emits pollutants, contaminating the air that we breathe and leading to adverse health effects. Major sources responsible for emitting pollutants into the atmosphere including power generation, transport (including petrol and diesel vehicles), residential energy use, agriculture and industry.

Historically, energy from fossil fuels has dominated power generation (Figure. 2), but as the cost of establishing and maintaining renewable sources of power (such as wind and solar) continues to fall, these options are now frequently less expensive than the fossil fuel alternative. Research by the International Renewable Energy Agency published in 2018 took into account the lifetime cost of electricity in its calculations of cost comparisons to generate power from renewable sources versus fossil fuels. Although in most parts of the world newly commissioned power plants that use renewable sources, such as wind and solar, will be cheaper or at a similar cost than from fossil fuels, including coal, oil and gas<sup>18</sup>, companies continue to push outdated technologies with the outcome that fossil fuels continue to dominate, creating air pollution when cleaner alternatives are readily available.

## INTRODUCTION

Figure 2: Global active power plant capacity added per year, still active in 2014. Fossil fuel use has dominated energy supplies for decades, but the use of renewable technologies has been rapidly expanding, particularly since the mid-2000s<sup>19</sup>.



### 1.3 Air pollution, health and cost

Air pollution affects physical and mental health because it contributes to acute and chronic diseases that can reduce quality of life. Evidence from public health studies suggests that exposure to an air pollutant or combination of air pollutants, such as PM<sub>2.5</sub>, NO<sub>2</sub> or ozone, is associated with increased incidence of diseases including ischaemic heart disease (IHD), chronic obstructive pulmonary disease (COPD), lung cancer, lower respiratory infections, premature birth (preterm birth), type II diabetes, stroke and asthma<sup>20,21,22,23</sup>. Health impacts from air pollution generate economic costs from the cost of treatment, management of health conditions, and from work absences.

The properties and effects of air pollution vary from country to country; different locations are affected by different pollutants, pollution sources and environmental conditions. Combined with differences in population and lifestyle, the health impacts from air pollution change significantly depending on the geographical location<sup>24,25,26</sup>. For example, a computer modelling study<sup>27</sup> looked at seven different sources of PM<sub>2.5</sub> and ozone air pollution: industry; land traffic; residential and commercial energy; biomass burning; power generation; agriculture; and natural. Using the model, the researchers calculated premature mortality resulting from air pollution generated by each of the seven sectors. Of premature deaths attributed to air pollution globally in 2010, almost one-third were attributable to exposure (while outdoors) to air pollution from residential and commercial energy<sup>28</sup>, which was the principal source of air pollution-related premature deaths in India and China Mainland. Globally, land traffic was attributable for 5% of air pollution-related premature deaths and power generation for 14%. Countries where air pollution from land traffic emissions were particularly high included the US, Germany, Russia,

Turkey and Japan. The contribution of power generation to premature mortality was particularly high in the US, Russia, Turkey, China Mainland and Japan<sup>29</sup>.

The World Health Organisation (WHO) has created guidelines that describe the level of air pollution above which there is strong evidence of negative health impacts<sup>30</sup>. These guidelines are derived from the latest available evidence on the health effects of ambient air pollutants and undergo regular review<sup>31</sup>. In 2019, around 91% of the global population lived in places where levels of air pollution exceeded the WHO guidelines<sup>32</sup>.

### 1.4 Air pollution and climate

The phaseout of fossil fuels and switch to renewable forms of energy is beneficial both for reducing air pollution and mitigating anthropogenic climate change<sup>33</sup>. Events that may be affected by climate change such as sandstorms, wildfires and heat waves can worsen air pollution by, for example, increasing the quantity of particulate matter in the air. Reducing the health burden attributed to air pollution while simultaneously reducing emissions of climate pollutants can be achieved by, for example, removing coal from the energy industry or limiting emissions from the transport sector<sup>34</sup>.

Air pollution and the climate crisis are clearly linked. Replacements for the coal, oil and gas that are currently used to generate power, for transport and domestic heating will need careful consideration to ensure that alternative combustion processes are avoided. For example, if future energy generation is obtained by burning biomass the likely scenario is increased emissions of harmful air pollutants such as PM<sub>2.5</sub><sup>35</sup>. The pathways chosen to meet greenhouse gas-reduction targets must champion renewable technology and resource efficiency without negatively affecting air quality.

## 2.0 THE ECONOMIC COST OF AIR POLLUTION FROM FOSSIL FUELS

### 2.1 Introduction

This section presents the first global assessment of the economic burden of health impacts from fossil fuel air pollution. Analysis commissioned by Greenpeace Southeast Asia and carried out by the Centre for Research on Energy and Clean Air (CREA) has estimated the global and national impact of air pollution from fossil fuels. This chapter presents the findings from the analysis. Air pollution from non-fossil fuel sources (see section 1.1.2) is not included in the analysis.

The CREA/Greenpeace analysis suggests that air pollution from burning fossil fuels costs an estimated 3.3% (95% confidence interval 2.4–4.7%) of global gross domestic product, equivalent to US\$8 billion per day (95% confidence interval \$5.5–11.0bn) and 12,000 (95% confidence interval 9,000–17,000) premature deaths every day.

The assessment incorporates recent research that quantifies the contribution of fossil fuels to global air pollution levels. It uses published global datasets describing surface level concentrations of PM<sub>2.5</sub>, ozone and NO<sub>2</sub> to perform a health impact assessment and subsequent cost calculation for the year 2018. Full details of the methodology are provided in Appendix 1.

The health and economic impacts included in the CREA/Greenpeace analysis only consider fossil fuel-related air pollution and are shown in Table 1. Only those impacts for which there is sufficiently robust data relating pollutant concentrations to population level health impacts are included. Therefore, these figures represent only a proportion of the total burden of all air pollution. Finally, because not all real world health impacts from fossil fuel air pollution are included, the analysis presented in this chapter is a conservative estimate of the global impact of fossil fuels on air pollution, health and the economy.

The health impacts are determined by combining pollutant concentration maps<sup>36,37</sup>, representing the year 2018, with country or region-level demographic data and health statistics. Concentration response functions for each pollutant are used to relate a pollutant concentration to the response or impact of that pollutant on a population. Published concentration-response functions, described in Appendix 1, were used to calculate the impact of the mapped fossil fuel air pollution at a population level for the calendar year 2018. Using research carried out by the World Health Organization's project called 'Health risks of air pollution in Europe' (HRAPIE)<sup>38</sup>, it is possible to estimate the number of working days lost through exposure to PM<sub>2.5</sub>, but statistics are not available to calculate the impact of other air pollutants on work absences. The concentration response functions used to calculate the incidence of the health and economic outcomes shown in Table 1.

Previous studies have estimated global and regional mortality and disease incidence rates resulting from exposure to air pollution and fossil fuel derived air pollution. The methodology applied in this work builds on recent scientific findings and models of the health risks of air pollution exposure. An updated model of mortality risks due to outdoor PM<sub>2.5</sub> pollution in 2018 found substantially higher risks than earlier studies and consequently revised the estimates of premature deaths upwards<sup>39</sup>. A new cohort study on ozone similarly led to a large increase in deaths attributed to ozone pollution in 2017<sup>40</sup>. For these reasons, the estimates of death and disease in our study exceed many earlier results, sometimes leading to the number of deaths attributed to fossil fuels exceeding the total number of deaths from air pollution reported in some earlier studies. Full details of the health and economic impact calculations used in the CREA/Greenpeace analysis are provided in Appendix 1.

Table 1: Impacts included in the analysis by pollutant type\*.

Pollutant	Impact of pollutant exposure	Outcome
NO <sub>2</sub>	Asthma	Asthma prevalence New asthma cases in Children
	Non-communicable diseases and lower respiratory infections	Premature deaths Years of life lost
Ozone	Chronic obstructive pulmonary disease	Premature deaths Years of life lost
	Asthma	Emergency room visits
PM <sub>2.5</sub>	Work absences	Work absences
	Chronic obstructive pulmonary disease	Premature deaths
	Diabetes Caused by Chronic Diseases (Years Lived with Disability)	
	Ischaemic heart disease	
	Lung cancer	
	Lower respiratory infections	
	Lower respiratory infections in children under 5	
	Other non-communicable diseases and lower respiratory infections	
	Stroke	Years of life lost
	Asthma	
	Preterm birth	
	Chronic obstructive pulmonary disease	
	Diabetes Caused by Chronic Diseases (Years Lived with Disability)	
Stroke		
Lower respiratory infections in children under 5		
Non-communicable diseases and lower respiratory infections		

\*Although many health impacts are linked to PM2.5, NO2 and ozone, only those health impacts for which a robust relationship exists between changes in pollutant concentration and the incidence of disease have been included in the study.

## 2.2 Health impacts and costs

### 2.2.1 Health

The CREA/Greenpeace analysis suggests that an estimated 3 million premature adult deaths each year are attributed to cardiovascular diseases, respiratory diseases and lung cancer through exposure to PM<sub>2.5</sub> air pollution from fossil fuels. An estimated 500,000 premature deaths from chronic diseases are attributed to fossil fuel-related NO<sub>2</sub> pollution and 1 million premature deaths are attributed to fossil fuel-related ozone pollution annually. Combined, total premature deaths per year attributable to fossil fuel-related air pollution is estimated at 4.5 million. In the MENA region, fossil fuels related air pollution caused around 65,000 premature deaths in 2018.

The data calculated for this report estimates that 40,000 children may die before their fifth birthday due to illnesses related to exposure to PM<sub>2.5</sub> from fossil fuels and shows that those deaths occur mainly in low-

income countries. The estimated number for the MENA is 865 children. The loss of a child is tragic and devastating to the families affected. In economic terms, infant mortality has a high fiscal cost to society because that child is prevented from contributing to society in adulthood (Table 2).

Millions of people around the world are living with asthma and other chronic respiratory diseases where exposure to pollution from fossil fuel combustion is a contributing factor. Chronic diseases can lead to substantial healthcare costs and prevent people from participating in the workforce. Exposure to PM<sub>2.5</sub> is the leading cause of health and economic impacts arising from air pollution and in 2018 contributed to a higher number of premature deaths than those we can attribute to exposure to NO<sub>2</sub> and ozone combined (Fig. 3).





Table 2: The estimated global impact of fossil fuel-related air pollution on selected health and economic outcomes by pollutant in 2018. Provided are the upper and lower bounds of a 95% confidence interval (low, high) and a central estimate.

Pollutant	Impact	Total Number			Total Cost (Million US Dollars)		
		Low	Central	High	Low	Central	High
NO <sub>2</sub>	Premature Deaths	300,000	500,000	1,100,000	-	-	-
	Years of Life Lost	4,900,000	8,900,000	19,400,000	185,000	335,000	732,000
	New Cases of Asthma in Children	1,800,000	4,000,000	5,200,000	-	-	-
	Number of Children Living with Asthma Due to Air Pollution	7,800,000	16,100,000	19,600,000	8,000	16,000	19,000
Ozone	Premature Deaths	600,000	1,000,000	1,400,000	-	-	-
	Years of Life Lost	9,700,000	15,400,000	21,800,000	247,000	379,000	523,000
	Asthma (Emergency Room Visits)	2,800,000	5,600,000	5,600,000	500	1,000	1,000
PM <sub>2.5</sub>	Premature Deaths	2,300,000	3,000,000	3,700,000	-	-	-
	Years of Life Lost	48,700,000	62,700,000	77,700,000	1,385,000	1,766,000	2,173,000
	Asthma (Emergency Room Visits)	1,800,000	2,700,000	3,800,000	200	350	500
	Preterm Births	1,000,000	2,000,000	2,100,000	47,000	91,000	96,000
	Work Absences	1,503,200,000	1,755,200,000	2,002,200,000	86,000	101,000	115,000
Total	Premature Deaths	3,200,000	4,500,000	6,200,000	-	-	-
	Years of Life Lost	63,300,000	87,000,000	118,900,000	1,817,000	2,480,000	3,428,000

\*Low, central and high estimates are provided representing a 95% confidence interval.

Figure 3: The estimated global number of premature deaths from exposure to fossil fuel-related air pollution in 2018.



2.2.2 Economic cost

Data generated by the CREA/Greenpeace model suggests that an estimated annual cost of US\$2.9 trillion (central estimate), equivalent to 3.3% of global GDP or US\$8 billion per day, is attributed to air pollution from fossil fuels (Table 3, Figure 4). These costs are the result of respiratory and non-communicable diseases made more likely by elevated pollution levels. An economic valuation of the years of life lost through premature death is also included. The impact of premature death can be quantified using a measure known as 'years of life lost'. The personal tragedy of a premature death also brings an economic cost through

lost contributions to society and economy. This means that when premature deaths occur, especially in children and younger people, the economic cost can be large.

Costs of US\$350 billion and US\$380 billion are attributed to NO<sub>2</sub> and ozone air pollution from fossil fuels respectively, each equivalent to 0.4% of global GDP.


PM<sub>2.5</sub> air pollution leads to the greatest health impact and the greatest financial cost of the three pollutants. PM<sub>2.5</sub> from fossil fuels is attributed to increased work absences, causing an estimated 1.8 billion (central estimate) days of work absences annually worldwide (Table 3).

Table 3: The estimated annual global cost of fossil fuel-related air pollution in 2018\*

Pollutant	Impact	Low	Central	High
NO <sub>2</sub>	Total Cost (Billion US\$)	192	351	750
	% GDP	0.2%	0.4%	0.9%
Ozone	Total Cost (Billion US\$)	248	380	524
	% GDP	0.30%	0.40%	0.60%
PM <sub>2.5</sub>	Total Cost (Trillion US\$)	1.6	2.2	2.7
	% GDP	1.8%	2.5%	3.1%
	Work Absences (Days)	1,503,200,000	1,755,200,000	2,002,200,000
Combined	Total Cost (Trillion US\$)	2.09	2.9	4.0
	% GDP	2.4%	3.3%	4.7%

\*Low, central and high estimates are provided representing a 95% confidence interval.

Figure 4: The estimated annual global cost of fossil fuel-related air pollution in 2018. Low, central and high estimates are provided, representing a 95% confidence interval.



### 2.3 Regional examples of the cost of air pollution

This section presents findings generated by the CREA/Greenpeace data in particular relevance to different global regions.

#### 2.3.1 Health

The health impact of fossil fuel air pollution in a country or region is determined by factors including the nature and distribution of pollution sources, the local environment, weather conditions, background rates of disease not related to air pollution, population size and population density, among others.

The data generated in the CREA/Greenpeace analysis includes the projected number of asthma-related emergency room (ER) visits attributable to PM<sub>2.5</sub> and ozone. Globally, the findings estimate 7.7 million (confidence 4.8-10.0 million) asthma-related ER visits annually. Of these, 37,000 (24,000-47,000) ER visits are in Russia; 62,000 (37,000-83,000) in South Africa; 127,590 (78,450-166,450) in Egypt; 196,000 (125,000-248,000) in the US; and 266,000 (166,000-340,000) in Indonesia, for example. A breakdown of cost and premature death data by location is provided in Appendix 2.

Approximately 2 million (1,032,000-2,093,000) preterm births worldwide are attributed to PM<sub>2.5</sub> exposure as a result of fossil fuel use. Of these, an estimated 350,000 (184,000-367,000) are in China Mainland; 14,000 (6,700-14,500) are in South Africa; 981,000 (517,000-1,031,000) are in India; and 11,000 (6,000-12,000) are in Thailand.


The incidence of stroke has been linked to PM<sub>2.5</sub> exposure and the CREA/Greenpeace data estimates that 600,000 (268,000-904,000) deaths annually can be attributed to stroke relating to fossil fuel derived PM<sub>2.5</sub> exposure.

Across the EU, around 400,000 annual premature deaths are attributed to exposure to air pollution from fossil fuel use. Of those deaths three-quarters are related to PM<sub>2.5</sub> exposure with the remainder being related to NO<sub>2</sub> and ozone exposure (Table 4, Figure 5).

Table 4: The estimated number of premature deaths in the European Union attributable to fossil fuel air pollution in 2018. Low, central and high estimates are provided representing a 95% confidence interval

EU	Premature deaths in 2018		
	Low	Central	High
Total	289,000	398,000	567,000
PM <sub>2.5</sub>	229,000	295,000	367,000
NO <sub>2</sub>	38,000	69,000	152,000
Ozone	22,000	34,000	48,000


Figure 5: Estimated premature deaths in the EU attributable to fossil fuel air pollution by pollutant (2018). Low, central and high estimates are provided representing a 95% confidence interval.



In East Asia the absolute number of early deaths attributed to air pollution is dominated by those in China Mainland. Despite rapid improvements in PM<sub>2.5</sub> air quality since 2015<sup>41</sup>, pollution continues to affect a large proportion of the population. Annually 1.8 million (1.3-2.5 million) premature deaths are projected across China Mainland. The estimated number of annual premature deaths relating to fossil fuel emissions is 40,000 (28,000-61,000) in South Korea, 16,000 (12,000-24,000) in Taiwan and 100,000 (75,000-150,000) in Japan.

Across East Asia, the majority of premature deaths are attributed to PM<sub>2.5</sub> exposure (Figure 6). Chronic obstructive pulmonary disorder (COPD) is a leading cause of early deaths. Data from the present analysis attribute 582,000 (366,000-827,000) deaths to COPD-related to PM<sub>2.5</sub> exposure from fossil fuels in China Mainland, approximately 40% of the global incidence. In South Korea 5000 (3,000-7,000) premature deaths from COPD are projected with 4,000 (2,000-6,000) in Taiwan; and 15,000 (9,000-22,000) in Japan.

Figure 6: Estimated premature deaths in East Asia attributable to fossil fuel air pollution by pollutant (2018). Low, central and high estimates are provided representing a 95% confidence interval.



Premature deaths attributed to fossil fuel air pollution in the Southeast Asian nations of Vietnam, Laos, Thailand, Myanmar, Singapore, Cambodia, Malaysia, Indonesia, Philippines, Brunei and Timor Leste are shown in Table 5 and Figure 7. The greatest number of premature


deaths are projected in Indonesia and Vietnam, where deaths from COPD attributable to PM<sub>2.5</sub> are estimated to be 17,000 (10,000-25,000) and 10,000 (5,000-17,000) respectively.

Table 5: Estimated premature deaths in Southeast Asia attributable to fossil fuel air pollution (2018). Low, central and high estimates are provided representing a 95% confidence interval.

Southeast Asia	Premature deaths in 2018		
	Low	Central	High
Vietnam	28,000	41,000	58,000
Laos	1,400	2,000	2,900
Thailand	17,000	24,000	34,000
Myanmar	12,000	18,000	24,000
Singapore	890	1,000	2,000
Cambodia	1,900	2,800	4,100
Malaysia	4,300	6,600	10,000
Indonesia	30,000	44,000	61,000
Philippines	11,000	17,000	27,000
Brunei Darussalam	20	30	40
Timor Leste	10	20	30



Figure 7: The estimated number of premature deaths in Southeast Asia attributable to fossil fuel air pollution by pollutant (2018). Low, central and high estimates are provided representing a 95% confidence interval.



In Middle Eastern and North African countries, the estimated annual number of premature deaths from fossil fuel air pollution is around 65,000. Egypt has the highest estimated number of premature deaths from fossil fuel air pollution with 32,000 premature deaths in 2018 (Table 6, Figure 8). Of those, 4,600 (2,300-7800) are related to COPD, 4,000 (1,500-7,000) are related to stroke, and 15,000 (12,000-18,000) are related to

ischaemic heart disease and PM2.5 exposure attributed to fossil fuel use. The estimated number of annual premature deaths attributable to fossil fuel air pollution for all countries analysed is provided in Appendix 2. Lebanon has the highest estimated premature death rate attributed to fossil fuels air pollution followed by Egypt and Syria (Table 8, Figure 9).

Table 6: The estimated number of premature deaths in the Middle East and North Africa attributable to fossil fuel air pollution (2018). Low, central and high estimates are provided representing a 95% confidence interval.

Middle East and North Africa	Premature deaths in 2018		
	Low	Central	High
Algeria	2,100	3,000	4,300
Bahrain	200	300	400
Egypt, Arab Rep.	22,000	32,000	51,000
Iraq	2,500	3,500	4,800
Jordan	800	1,200	1,900
Kuwait	290	410	600
Lebanon	1,800	2,700	4,200
Libya	600	900	1,300
Morocco	3,300	5,100	7,500
Oman	140	210	300
Palestine	400	500	700
Qatar	140	230	410
Saudi Arabia	2,200	3,300	5,000
Syrian Arab Republic	3,100	4,700	7,100
Tunisia	1,300	2,100	3,100
United Arab Emirates	900	1,500	2,400
Yemen, Rep.	1,800	3,100	5,200

Figure 8: The estimated number of premature deaths in the Middle East and North Africa attributable to fossil fuel air pollution by pollutant (2018). Low, central and high estimates are provided representing a 95% confidence interval.




Table 7: The estimated number of premature deaths in the MENA region attributable to fossil fuel air pollution in 2018. Low, central and high estimates are provided representing a 95% confidence interval.


MENA	Premature deaths in 2018		
	Low	Central	High
Total	43,000	65,000	100,000
PM <sub>2.5</sub>	32,000	44,000	59,000
NO <sub>2</sub>	5,000	10,000	24,000
Ozone	6,000	11,000	17,000

Table 8: Estimated premature death rate per 1000 people attributed to fossil fuel related air pollution per country in the MENA region 2018. Low, central and high estimates are provided representing a 95% confidence interval.

Country	Premature deaths in 2018		
	Low	Central	High
Algeria	0.05	0.07	0.10
Bahrain	0.12	0.17	0.27
Egypt	0.22	0.33	0.51
Iraq	0.7	0.09	0.13
Jordan	0.8	0.13	0.19
Kuwait	0.7	0.10	0.15
Lebanon	0.26	0.39	0.61
Libya	0.9	0.13	0.19
Morocco	0.9	0.14	0.21
Oman	0.3	0.04	0.06
Palestine	0.8	0.11	0.15
Qatar	0.5	0.08	0.15
Saudi Arabia	0.7	0.10	0.15
Syria	0.19	0.28	0.42
Tunisia	0.11	0.18	0.27
United Arab Emirates	0.09	0.15	0.25
Yemen, Rep.	0.06	0.11	0.18

**THE ECONOMIC COST OF AIR POLLUTION FROM FOSSIL FUELS**

**Figure 9: Total premature deaths and premature death rate attributed to fossil fuel related air pollution per country in the MENA region (Central Estimate).**



**2.3.2 Economic cost**

Population size is a significant driver of the absolute cost of air pollution for society. Also important are factors such as the local availability and cost of healthcare and the rate of infant mortality. Nations that have large populations often have a large absolute cost burden from fossil fuel-related air pollution, and where high air pollution levels intersect with dense populations the impact is magnified.

The cost of fossil fuel air pollution equates to a large percentage of many nations' GDP. The projected cost of fossil fuel air pollution as a percentage of GDP is greatest in China Mainland, where it equates to 6.6% (confidence 4.7-9.0%) of GDP. By comparison, it is 3.4% (2.2-5.1%) and 2.5% (1.8-3.7%) in South Korea and Japan respectively. In Bulgaria, Hungary, Ukraine, Serbia, Belarus, India, Romania and Bangladesh the cost (central estimate) is greater than 5% of GDP. The estimated cost of fossil fuel-

related air pollution in the Middle East and North Africa is shown in Table 7 and Table 8.

Across East Asia the estimated cost of fossil fuel-related air pollution is high, accounting for more than 2% of GDP across East Asia. Work absences in 2018 from exposure to fossil fuel-related PM2.5 are estimated at 748 million (642-853 million) days in China Mainland, costing the economy an estimated US\$39 billion (US\$33-US\$44 billion). In South Korea, Taiwan and Japan the figure is 18 million (16-21 million) days, 5 million (4-6 million) days and 20 million (17-23 million) days, respectively.

In Southeast Asia, the estimated cost of fossil fuel-related air pollution is greater than 2% of GDP in Vietnam, Laos, Thailand and Myanmar (central estimate). Only in Brunei and Timor Leste is the figure less than 1% of GDP (Table 9).

**Table 9: The estimated annual cost (% GDP) of fossil fuel-related air pollution in Southeast Asia in 2018. Low, central and high estimates are provided representing a 95% confidence interval.**

Southeast Asia	% GDP total		
	Low	Central	High
Vietnam	1.8%	2.8%	4.0%
Laos	1.8%	2.9%	4.1%
Thailand	1.4%	2.1%	2.9%
Myanmar	1.8%	2.7%	3.6%
Singapore	0.7%	1.1%	1.8%
Cambodia	1.0%	1.5%	2.1%
Malaysia	0.8%	1.3%	1.9%
Indonesia	0.8%	1.1%	1.6%
Philippines	0.8%	1.2%	1.9%
Brunei Darussalam	0.3%	0.4%	0.6%
Timor Leste	0.1%	0.1%	0.2%

**THE ECONOMIC COST OF AIR POLLUTION FROM FOSSIL FUELS**

The estimated cost of fossil fuel-related air pollution as a percentage of GDP in the Middle East and North Africa is greatest in Egypt, Lebanon, Bahrain and the UAE (Table 10). Work absences resulting from exposure to fossil fuel-related PM2.5 reach 15 million (13-17 million) days in Egypt; 1.3 million (1.1-1.5 million) days in Lebanon; 460,000 (400,000-530,000) days in Bahrain; and 2.7 million (2.3-3.1 million)

days in the UAE. Egypt, Saudi Arabia and the United Arab Emirates bear the highest costs from fossil fuel air pollution in the MENA at an estimated US\$6.9 billion, US\$6 billion and US\$5.9 billion per year, respectively (Table 11).

The total cost of fossil fuel air pollution for all countries analysed is provided in Appendix 2.

**Table 10: The estimated annual cost (% GDP) of fossil fuel-related air pollution in the Middle East and North Africa in 2018. Low, central and high estimates are provided representing a 95% confidence interval.**

Middle East and North Africa	% GDP total		
	Low	Central	High
Algeria	0.3%	0.5%	0.7%
Bahrain	0.9%	1.4%	2.1%
Egypt, Arab Rep.	1.8%	2.8%	4.2%
Iraq	0.5%	0.8%	1.1%
Jordan	0.7%	1.1%	1.5%
Kuwait	0.6%	0.9%	1.2%
Lebanon	1.3%	2.0%	3.0%
Libya	0.6%	0.9%	1.3%
Morocco	0.6%	0.9%	1.4%
Oman	0.3%	0.4%	0.6%
Palestine	0.5%	0.8%	1.0%
Qatar	0.5%	0.8%	1.3%
Saudi Arabia	0.5%	0.7%	1.1%
Syrian Arab Republic	No Data	No Data	No Data
Tunisia	0.6%	1.0%	1.5%
United Arab Emirates	0.8%	1.4%	2.2%
Yemen, Rep.	0.5%	1.0%	1.6%

**Table 11: Estimated total cost of fossil fuel related air pollution per country in the MENA region. Low, central and high estimates are provided representing a 95% confidence interval.**

Country/Region	Estimated total cost (Million USD)		
	Low	Central	High
Algeria	530	840	1,100
Bahrain	330	510	750
Egypt, Arab Rep.	4,400	6,900	10,000
Iraq	1,400	2,100	2,800
Jordan	300	490	700
Kuwait	840	1,300	1,700
Lebanon	890	1,400	2,100
Libya	300	470	660
Morocco	670	1,100	1,600
Oman	200	320	430
Palestine	80	120	160
Qatar	1,000	1,600	2,400
Saudi Arabia	3,800	6,000	8,800
Syrian Arab Republic	No Data	No Data	No Data
Tunisia	240	400	590
United Arab Emirates	3,500	5,900	9,400
Yemen, Rep.	150	280	450

## 3.0 WHAT CAN BE DONE ABOUT ANTHROPOGENIC AIR POLLUTION?



Solar rooftop at Luang Suan Hospital in Thailand  
©Arnaud Vittet/Greenpeace



Salt fields near a wind farm in Guimaras, Philippines  
© Veejay Villafranca/Greenpeace

### 3.1 Case study one: Switch to sustainable transport

The ways in which humans travel — particularly in highly populated urban areas — must change if we are to tackle the dual threats of air pollution and climate emergency<sup>46,47</sup>. As millions of people take private vehicles on a daily basis for work, school or leisure, neighborhood streets are not only clogged with traffic but diesel and petrol engines contribute to poor air quality and lead to an increased concentration of atmospheric greenhouse gases. It is clear that to reduce harmful levels of particulates and halt global climate heating, a transport revolution is needed, one that ensures a clean, carbon-neutral and universal mobility system for all. Our cities need to support lifestyles that are healthy for residents and for the planet. Low-cost, active and carbon neutral transportation is an important part of this transition, having the combined benefits of reducing urban pollution, greenhouse gas emissions, and rates of cardiovascular disease, cancer, obesity, diabetes, mental illness, and respiratory disease<sup>48</sup>. One of the most important steps governments can take to initiate a move towards a sustainable transport system is to set a phase-out date for diesel and petrol cars, while implementing various urban transport measures, such as restricting cars' access into certain neighborhoods or districts, banning whole categories of cars within city limits, and promoting car-free days. Such initiatives allow residents to imagine what their city could be like without congestion and particulate pollution, with the added benefit of encouraging physical activity. In this case study, we investigate initiatives that national governments have undertaken to improve public health and air quality by removing petrol and diesel cars from city streets.

There are many examples of city authorities that have undertaken initiatives to create cleaner air, for example by promoting pedestrian and cycle-friendly spaces and alternatives to private vehicle transport, such as car clubs or vehicle sharing schemes, and public transport powered by renewable energy. In the UK, Transport for London announced that (correct as of January 2020), four bus routes through the city centre are fully electric<sup>49</sup>. In Shenzhen, all diesel public buses were replaced with electric vehicles in 2018, making it the world's first fully electric fleet<sup>50</sup>. In the United States, New York City's Metropolitan Transportation Authority (MTA) is working towards a zero-emission bus fleet and is in the final stages of a three-year trial of all-electric buses<sup>51</sup>. In 2019, Oslo closed central city streets to privately owned vehicles and cut city centre parking spots<sup>52</sup>. Oslo residents can enjoy a cleaner environment, where walking, cycling, and public transport use is encouraged. Restricting private vehicle use is an effective strategy to reduce fossil fuel use and improve

**Realistic scenarios for phasing out fossil fuels are capable of simultaneously reducing air pollution and greenhouse gas emissions<sup>42</sup>. Decarbonising the global economy can bring rapid health gains for society, especially by reducing exposure to air pollutants, most notably PM<sub>2.5</sub> which has the greatest impact on our health<sup>43</sup>.**

Solutions to the air pollution crisis, such as emissions controls in Europe, have helped bring about huge reductions in health impacts for citizens. Policy to reduce air pollution does not need to be expensive, and, where interventions are costly to implement, the benefits can outweigh the costs. For example, the United States Clean Air Act has returned substantially greater economic benefits in relation to the costs of implementation, and the benefits exceeded a ratio of 30:1 over the period 1990–2020. Put another way, for every US\$1 invested, the US economy saw benefits of at least US\$30 returned<sup>44</sup>. Many strategies to control pollution are likely to be cost-effective in cities and countries whatever their level of income<sup>45</sup>.

In this section we demonstrate with examples that it is possible and affordable to drastically reduce the health burden of air pollution from fossil fuels, through actions that also support efforts to mitigate anthropogenic climate change. This section focuses on two different fossil fuel burning sectors: transport and oil; and electricity and coal. We demonstrate that action on air pollution is practical, feasible and cost effective.

air quality, and in conjunction with alternate and public transport systems, access can be maintained for those unable to walk and cycle<sup>53</sup>.

No city has become entirely vehicle free, but car-free days which have operated in cities for many years provide an opportunity to see the potential for health and environmental co-benefits. Car-free events take place in cities around the world, often annually but in some locations, like Bogota, Colombia, roads are closed to cars on a weekly basis<sup>54</sup>. These examples show that, although occasional car-free days are a positive first step, to be effective for human health, car-free events should take place on a regular or continuous basis and attract a high proportion of the community.

As we can learn from the car-free cases, ensuring streets aren't clogged with cars leads to significant health and financial benefits. However, car-free cases are only a partial policy approach to resolve congestion and air pollution in urban areas. The bigger shift is happening globally to move the transport system away from fossil fuels and approach mobility in a way that is sustainable and equitable for all.

More than 15 countries have announced plans to phase out new petrol and diesel cars, and, in some cases, hybrids. Although not perfect, these announcements send a strong signal to markets that there is no future for fossil fuel vehicles. Both to ensure safe air and avoid the worst impacts of climate change, it is necessary to phase out the internal combustion engine and develop alternatives to mass car ownership<sup>55</sup>.

It is clear that with technological and social changes, cities can eliminate fossil fuel-powered vehicles, rapidly reduce pollution and help find solutions to climate change. Major cities can pioneer this change, leading the charge toward more sustainable urban spaces that put social equity and justice at the heart of creating a carbon-neutral mobility system for all. In turn this will shape national and global debates on the future of urban transport<sup>56</sup> and empower national governments to set ambitious phase-out dates for petrol and diesel cars while investing in public transport and e-mobility.

One of the longest-running examples of a car-free day event in a city is the Ciclovía in Bogotá, Colombia which was initiated in 1974. The Ciclovía is a weekly event that takes place every Sunday and public holiday in central Bogotá, in which roads are closed to motorised traffic to allow cyclists, skaters and pedestrians traffic-free access to 120 kilometres (74.6 miles) of roads. Similar events take place in more than 15 countries in the Americas and the Caribbean. The aim of the Ciclovía was to encourage more people to take up physical activity, and some people may choose to exercise during Ciclovía precisely because there is no traffic and lower air pollution<sup>57</sup>. The inclusive nature of the Ciclovía boosts its potential to promote activity in children and prevent them from developing obesity<sup>58</sup>.

#### **The cost-benefit ratio in terms of health cost savings of the Bogotá Ciclovía**

In terms of health costs savings, the estimated cost-benefit ratio of the Bogotá Ciclovía ranges from US\$3.20 to US\$4.30 per US\$1 invested in the programme<sup>59</sup>. The range reflects uncertainty in the number of adult users, which in a study period between 2005 and 2009, was estimated to be between 516,600 and 1,205,635. A cost-benefit ratio calculated for Medellín, Colombia's second city, which runs a similar Ciclovía programme, found that every US\$1 invested yielded a health cost saving of US\$1.80. The total benefits of the Ciclovía events however may have been underestimated because only some beneficial factors were included in the analysis. The cost of the Ciclovía programme, based on data collected in the study period in 2005 and 2009, was estimated to be US\$6 per capita. For open streets events to be successful, the researchers suggest that the route should pass through different neighbourhoods, be promoted among under-represented ethnic and age groups, and have secure funding to ensure longevity<sup>60</sup>.

### **3.2 Case study two: Generate electricity from renewables, not fossil fuels**

Modelling studies suggest that 65% of air pollution related global premature mortality can be attributed to fossil fuel emissions<sup>61</sup>. Replacing fossil fuels with renewable energy from sources such as wind and solar reduces both greenhouse gas emissions and emissions of air pollutants, creating a dual benefit for climate and human health. This transition is both feasible and achievable, and power generated by renewable systems is increasing used globally, as technology has matured and installation costs have fallen drastically.

In the US between 2007 and 2015, the combined contribution from wind and solar to the national grid, as well as from distributed photovoltaic power sources, increased from ~10GW to ~100GW<sup>62</sup>. Pollutant emissions avoided by this wind power generation are estimated to have delivered air quality and public health savings of between US\$28.4-107.9 billion through the avoidance of an estimated 2,900-12,200 premature deaths, as well as climate benefits savings of US\$4.9-98.5 billion. The estimated benefits due to solar power generation were US\$1.3-4.9 billion for air quality and public health and US\$0.4-8.3 billion for climate benefits, with an avoided 100-500 early deaths.

Air pollution benefits from the closure of coal-fired power stations can exceed the value of the electricity they generated<sup>63</sup>. As we adopt renewable power and phase out fossil fuels, large health benefits can be realised by prioritising infrastructure with the greatest potential to reduce exposure to air pollutants. Power generation from coal, oil and gas contribute to air pollution and the climate crisis. Phasing out these technologies promises environmental and health benefits. The benefits of a fossil fuel phaseout can be most significant for coal power stations because of the higher SO<sub>2</sub>, NO<sub>x</sub>, and PM<sub>2.5</sub> emission rates associated with coal<sup>64</sup>. Emission standards and plant closures in the United States power generation industry drove emission reductions of 20%, 72%, 50% and 46% for CO<sub>2</sub>, SO<sub>2</sub>, NO<sub>x</sub> and PM<sub>2.5</sub> respectively between 2007 and 2015.

#### **Examples of health benefits that have been measured from observations of closures of coal-fired power plants.**

The closure of existing coal-fired power plants is beneficial to the environment and human health. Air quality studies show that the closure of coal-fired power plants brings health benefits to people living and working in their vicinity.

- One study that evaluated the impact of coal-fired power plants on childhood development<sup>65</sup> followed two groups of non-smoking pregnant women living within 2.5 kilometers of the Tongliang power station in Chongqing, China. The first group (n=150) was enrolled in 2002 and had been exposed to polyaromatic hydrocarbons from a power plant; the second group (n=158) were enrolled in 2005 and had not been exposed to the same source because the plant closed in 2004. Exposure to PAHs has been linked to developmental problems, particularly in unborn and young children. The coal power plant which operated between December and May until 2004 did not use modern pollution reduction technology. The study found that reduced exposure to PAH in the second group was associated with beneficial effects on neurodevelopment as well as molecular changes related to improved brain development and health. The researchers recommend reducing exposure to toxic pollutants to help enhance neurodevelopment.

- Monitoring<sup>66</sup> that took place from 2011 to 2014 in Pittsburgh, Pennsylvania, United States, found that, following the closure of three coal-fired power plants, there was a downward trend in ambient PM<sub>2.5</sub> levels. The researchers used PM<sub>2.5</sub> measurements from 12 ground stations and from satellites to monitor the aerosol optical depth (AOD), observing a decline in AOD over the study period.

- A well-documented case study<sup>67</sup> in Dublin, Ireland, describes how a 1990 ban on bituminous coal sales reduced wintertime black smoke by 70% and decreased deaths from respiratory illness by 15% (about 116 people per year).



Air pollution protest in Bangkok  
©Wason Wanichakorn/Greenpeace



©TUPUNGATO / SHUTTERSTOCK.COM

## 4.0 CONCLUSIONS

### 4.1 Costs

We estimate that air pollution resulting from fossil fuels causes approximately 4.5 million premature deaths each year globally. Air pollution from fossil fuels costs us an estimated 1.8 billion lost working days worldwide per year through poor health. This hits our economy and seriously damages people's well-being. Combining health costs and work absences, an economic cost equivalent to 3.1% of global gross domestic product, or US\$8 billion per day, is estimated to arise from fossil fuel-related air pollution, according to this study.

The air pollution health crisis is driven by burning fossil fuels, which – because burning coal, gas and oil emits greenhouse gases – also contributes to the climate emergency. Governments making investments to replace fossil fuels with clean renewable energy stand to benefit from long-term economic returns and to deliver improved health and wellbeing to their citizens.

The need to move away from reliance on fossil fuels is clear and, while the cost of our reliance on coal, oil and gas continues to soar, life-saving alternatives are increasingly widespread and affordable.

### 4.2 Transport

Car-free initiatives have demonstrated that radical changes to our transport systems have the potential to boost physical activity, reduce emissions of harmful air pollutants and greenhouse gases and improve health. The money saved through these health benefits has been demonstrated to return multiple times the cost of implementation. It also shows the extent to which fossil fuel companies are currently profiting at the expense of our communities, who pay the price for air pollution. Investment in alternative sustainable alternative transport systems can be financially sound and positive for health, wellbeing and the planet.

Our transport systems urgently need to be reorganised so that they use energy and resources efficiently and operate without either directly or indirectly emitting harmful pollutants. Our cities need fewer and cleaner vehicles operating alongside greater use of public transport and widespread investments in shared mobility, walking and cycling. To achieve this, national governments must implement ambitious phase-out dates for diesel and petrol cars while investing in sustainable transport and enabling low-carbon alternatives, such as walking or cycling. Pioneering cities and regions can lead the way, helping to shape national and global debates on the future of urban transport.

### 4.3 Energy

A phaseout of existing fossil fuel power plants and an end to the construction of new projects is essential to limit global warming to 1.5°C above pre-industrial levels, but it will also reduce the emission of air pollutants that today disperse over hundreds of kilometers.

Emissions from coal combustion have been linked to a broad range of illnesses, including developmental problems in children and premature deaths from respiratory illnesses. Deployment of renewable technology in the electricity grid in parts of the US has reduced reliance on fossil fuels and reduced the emission of pollutants. The removal of fossil fuels is helping to prevent premature deaths and bring vast savings in health costs.

Economic savings from the improved air quality resulting from coal-fired power station closures can exceed the value of the electricity they generated<sup>68</sup>. Moving our energy generation sector from fossil fuels to renewables is an essential step towards preventing catastrophic climate change and protecting our health. A just transition to renewable energy is feasible, and we can't afford to wait any longer. Cities, governments and companies need to take action now.

# GLOSSARY

<b>Air pollutant</b>	An unwanted substance found in the air in the form of a solid particle, a liquid droplet or a gas. The substance may be harmful to human health, or damaging to the environment.
<b>ALRI</b>	Acute lower respiratory illness.
<b>AOD</b>	Aerosol optical depth. Aerosols are solid particles in the atmosphere that may be of different sizes and from natural or anthropogenic sources. Aerosols can have different effects depending on what they are made from and the geographical location. Measuring aerosol optical depth can indicate the quantity of aerosol depending upon how much light passes through.
<b>Central estimate</b>	See: Confidence interval.
<b>CEV</b>	Cerebrovascular disease.
<b>Confidence interval</b>	Scientific studies that use computer models do not give results with absolute certainty. Instead a range is provided (known as an 'interval'). The 'confidence range' is the range that is most likely to contain the true value. A 95% confidence interval means that with 95% probability, reality is somewhere inside the confidence interval, and a 5% chance it is outside the interval (either higher or lower than the range of numbers in the range). The value with the highest probability to be the true value is called the central estimate. It is somewhere inside the confidence interval. The bounds of the confidence interval are called the low and the high estimate.
<b>COPD</b>	Chronic obstructive pulmonary disease.
<b>DEFRA</b>	The Department for Environment, Farming and Rural Affairs of the government of the United Kingdom.
<b>Dust</b>	Solid airborne particles. A subclass of dust is PM2.5.
<b>GDP</b>	Gross domestic product.
<b>GNI</b>	Gross national income.
<b>High estimate</b>	See: Confidence interval.
<b>IHD</b>	Ischaemic heart disease.
<b>Low estimate</b>	See: Confidence interval.
<b>MODIS</b>	Moderate resolution imaging spectroradiometer.
<b>NO<sub>2</sub></b>	Nitrogen dioxide.
<b>NO<sub>x</sub></b>	Nitrogen oxides. A generic term for NO and NO <sub>2</sub> , a group of trace gases that are harmful to human health.
<b>O<sub>3</sub></b>	Ozone.
<b>PM<sub>2.5</sub></b>	Fine particulate matter.
<b>ppb</b>	Parts per billion. The number of units of mass of a contaminant per 1000 million units of total mass.
<b>PPP</b>	Purchasing power parity is a currency exchange rate used to determine the value of an international dollar such that it has the same purchasing power over gross national income as a United States dollar has in the United States.
<b>SO<sub>2</sub></b>	Sulfur dioxide.
<b>WHO</b>	World Health Organization
<b>µg/m<sup>3</sup></b>	Microgram per cubic meter. The mass of a substance in milligrams, in one cubic metre of a gas.

# APPENDIX 1: METHODOLOGY

This paper presents the first global estimate of the economic burden caused by air pollution from fossil fuels. We use global datasets describing surface level concentrations of PM<sub>2.5</sub>, ozone and NO<sub>2</sub> to perform a health impacts assessment and subsequent cost calculation. The health impacts are determined by combining pollutant concentration maps with population data, country or region-level health statistics and pollution exposure response functions for the year 2018. The complete methodology is presented below.

## Exposure to air pollution from fossil fuels

The concentrations of PM<sub>2.5</sub> and ozone linked to fossil fuel emissions have previously been estimated globally for 2015<sup>69</sup>, and total NO<sub>2</sub> pollution levels for the year 2011 have been mapped at high resolution<sup>70</sup>. As NO<sub>x</sub> emissions sources are heavily dominated by fossil fuel burning, we assume that NO<sub>2</sub> concentrations above the thresholds described in Table A1 represent NO<sub>2</sub> resulting from fossil fuel use. For example, for deaths linked to NO<sub>2</sub> exposure, the threshold is a concentration 20 µm/m<sup>3</sup> to link those deaths to fossil fuels. This assumption can be considered conservative because NO<sub>x</sub> emissions are heavily dominated by fossil fuels globally, particularly in cities where the majority of harmful exposure takes place.

The surface PM<sub>2.5</sub> and NO<sub>2</sub> data then are scaled to represent concentrations in 2018. The scaling is completed using satellite based observations of the respective years using MODIS and OMI data products. NASA MODIS (Moderate Resolution Imaging Spectroradiometer) is an Earth observation instrument that is carried on two NASA satellites, Terra and Aqua. It monitors aerosols in the atmosphere and completes a scan of the globe every 1-2 days<sup>71</sup>. OMI (Ozone Monitoring Instrument) monitors pollutants including NO<sub>2</sub> from the NASA AURA satellite. These satellite-based measurements quantify the total atmospheric amount of each pollutant, rather than surface concentrations, and are only used to adjust the surface level maps. Ozone is not adjusted from 2015 levels.

To calculate pollutant exposure, ozone and the adjusted 2018 PM<sub>2.5</sub> and NO<sub>2</sub> data are combined with population and health datasets updated to the latest available data (2017-2018)<sup>72,73,74,75</sup>.

## Health impacts and costs

We use the adjusted 2018 PM<sub>2.5</sub>, ozone and NO<sub>2</sub> data and concentration-response functions to determine the incidence of health impacts for a given population<sup>76</sup>. A concentration response function relates a pollutant concentration to the response or impact of that pollutant on a population. Deaths, years of life lost and years lived with disability due to PM<sub>2.5</sub> exposure are calculated using non-linear, age-specific risk derived by Burnett et al<sup>77</sup> and Global Burden of Disease, which give the increase in risk of different health effects as a

function of pollutant concentration, compared with clean air. Other health impacts are projected using log-linear risk functions that are expressed as increase in relative risk per 10µm/m<sup>3</sup>, or 10 ppb increase in pollutant concentration, and an assumed no-risk threshold, usually based on lowest concentration in study datasets that detected health risks. The following health impacts are considered:

### Deaths of small children from lower respiratory infections

This health impact is assessed using the PM<sub>2.5</sub> concentration results from Lelieveld et al (2019) and the Global Burden of Disease risk function for lower respiratory diseases<sup>78</sup>.

The economic losses from these air pollution-related deaths are assessed based on the resulting reduction in life expectancy, with one year of life lost equating to economic losses of EUR56,000 in the European Union, following the EEA cost-benefit methodology<sup>79</sup>, and adjusted by GNI PPP by country or region, with an elasticity of 0.9 as recommended by OECD<sup>80</sup>.

### Diabetes, asthma and other chronic respiratory diseases, and disabilities caused by stroke

The Global Burden of Disease project<sup>81</sup> has quantified the degree of disability caused by each disease into a "disability weight" that can be used to compare the costs of different illnesses.

The economic cost of disability and reduced quality of life caused by diabetes and chronic bronchitis is assessed based on these disability weights, combined with the economic valuation of disability used by the UK environmental regulator DEFRA<sup>82</sup>. The valuation is adjusted by GNI PPP for other countries. For example, type 2 diabetes without complications has a disability weight of 4.9%, meaning that the cost of one year lived with diabetes is estimated at 4.9% of the cost of one year lived with disability, or US\$4000 in the UK and US\$1600 at world average income level.

The economic cost of asthma related to fossil fuel pollution is assessed based on two indicators: new cases of asthma linked to NO<sub>2</sub> exposure, and emergency room visits related to PM<sub>2.5</sub> and ozone exposure. An assessment of the direct and indirect cost per year associated with childhood asthma, including medical costs and loss of income to the child's caregiver, found a cost of US\$3,800 and US\$4,000 in two different communities in California, US<sup>83</sup>. The midpoint of these two valuations is used for the estimates, adjusted by the ratio of California's Gross Regional Product to US national average, and by GNI PPP for other countries. The cost of an emergency room visit is taken from the same study.

Exposure to PM<sub>2.5</sub> is very likely linked to an even larger number of new asthma cases globally than exposure to NO<sub>2</sub>, but uncertainty in the estimates is large<sup>84</sup>, so this

**APPENDIX 1: METHODOLOGY**

effect is not included. Instead, we include the economic cost of emergency room visits for asthma linked to PM<sub>2.5</sub> and ozone exposure, which is only a small part of the overall cost of the burden of asthma linked to PM<sub>2.5</sub>.

**Preterm Birth**

PM<sub>2.5</sub> exposure to pregnant women increases the likelihood of preterm birth and low birth weight, which in turn increases the risk of many health and development issues throughout a baby's life. Approximately 2 million preterm births per year can be attributed to the exposure of pregnant women to PM<sub>2.5</sub> pollution from fossil fuels specifically, based on CREA analysis using the Lelieveld et al (2019) concentration results and the concentration-response relationship established by Trasande et al<sup>85</sup>. The same study estimated the economic

costs of a preterm birth, primarily lower economic productivity and increased health care costs, at US\$300,000 per birth in the US. This concentration-response function and cost estimate is used in the analysis; the valuations are adjusted using GDP PPP within regional groups.

**Work Absence**

Exposure to PM<sub>2.5</sub> air pollution from fossil fuels leads to increased work absences due to illness (sick leave). We estimate the incidence of work absences using WHO recommended concentration response functions<sup>86</sup>. The economic cost of these sick leaves is evaluated at EUR130 per day in the European Union, based on EEA recommendations<sup>87</sup>, and adjusted for other countries based on GDP PPP.

Table A1. Concentration Response Functions\*

Concentration Response Functions							
Health effects	Exposure	Risk ratio	Concentration change	No-risk threshold	Unit	Reference	Incidence data
Asthma emergency room visits, children	PM <sub>2.5</sub>	1.025 (1.013,1.037)	10	6	µg/m <sup>3</sup>	Zheng 2015	Anenberg et al 2018
Asthma emergency room visits, children	O <sub>3</sub>	1.018 (1.01,1.024)	10	2	ppb	Zheng 2015	Anenberg et al 2018
Asthma emergency room visits, adults	PM <sub>2.5</sub>	1.023 (1.015,1.031)	10	6	µg/m <sup>3</sup>	Zheng 2015	Anenberg et al 2018
Asthma emergency room visits, adults	O <sub>3</sub>	1.018 (1.012,1.022)	10	2	ppb	Zheng 2015	Anenberg et al 2018
Preterm births	PM <sub>2.5</sub>	1.15 (1.07, 1.16)	10	8.8	µg/m <sup>3</sup>	Trasande et al 2016	Chawanpaiboon et al 2019
Deaths, chronic obstructive pulmonary disease	O <sub>3</sub>	1.12 (1.08,1.16)	10	35	ppb	Malley et al 2017	GBD 2017
Deaths, non communicable diseases and lower respiratory infections, adults	NO <sub>2</sub>	1.037 (1.021,1.080)	10	20	µg/m <sup>3</sup>	WHO HRAPIE 2013	GBD 2017
Years of life lost, non-communicable diseases and lower respiratory infections, adults	NO <sub>2</sub>	1.037 (1.021,1.080)	10	20	µg/m <sup>3</sup>	WHO HRAPIE 2013	GBD 2017
Work absences, days	PM <sub>2.5</sub>	1.046 (1.039,1.053)	10	0	µg/m <sup>3</sup>	WHO HRAPIE 2013	EEA 2014
New cases of asthma in children	NO <sub>2</sub>	1.26 (1.10,1.37)	10	2	ppb	Achakulwisut et al 2019	GBD 2017

\*Table references 88,89,90,91,92,93,94,95,96,97,98.

# APPENDIX 2.

## SUMMARY OF COST AND MORTALITY DATA

Impact attributed to fossil fuel-related air pollution by country in MENA						
Country/Region	Estimated total cost (Million USD)			Estimated total premature deaths (2018)		
	Low	Central	High	Low	Central	High
<b>Algeria</b>	530	840	1,100	2,100	3,000	4,300
<b>Bahrain</b>	330	510	750	200	300	400
<b>Egypt, Arab Rep.</b>	4,400	6,900	10,000	22,000	32,000	51,000
<b>Iraq</b>	1,400	2,100	2,800	2,500	3,500	4,800
<b>Jordan</b>	300	490	700	800	1,200	1,900
<b>Kuwait</b>	840	1,300	1,700	290	410	600
<b>Lebanon</b>	890	1,400	2,100	1,800	2,700	4,200
<b>Libya</b>	300	470	660	600	900	1,300
<b>Morocco</b>	670	1,100	1,600	3,300	5,100	7,500
<b>Oman</b>	200	320	430	140	210	300
<b>Palestine</b>	82	120	160	400	500	700
<b>Qatar</b>	1,000	1,600	2,400	140	230	410
<b>Saudi Arabia</b>	3,800	6,000	8,800	2,200	3,300	5,000
<b>Syrian Arab Republic</b>	No Data	No Data	No Data	3,100	4,700	7,100
<b>Tunisia</b>	240	400	590	1,300	2,100	3,100
<b>United Arab Emirates</b>	3,500	5,900	9,400	900	1,500	2,400
<b>Yemen, Rep.</b>	150	280	450	1,800	3,100	5,200

Impact attributed to fossil fuel-related air pollution by country/region globally						
Country/Region	Estimated total cost (Million USD)			Estimated total premature deaths (2018)		
	Low	Central	High	Low	Central	High
<b>Afghanistan</b>	170	270	380	2,600	3,900	5,900
<b>Albania</b>	260	400	590	1,000	1,500	2,200
<b>Algeria</b>	530	840	1,100	2,100	3,000	4,300
<b>American Samoa</b>	No Data	No Data	No Data	0	0	1
<b>Andorra</b>	No Data	No Data	No Data	20	30	50
<b>Angola</b>	98	170	270	250	410	700
<b>Antigua and Barbuda</b>	3	5	7	4	6	7
<b>Argentina</b>	2,600	4,400	7,900	5,300	8,600	15,000
<b>Armenia</b>	270	370	480	1,300	1,700	2,200
<b>Australia</b>	3,900	6,100	8,900	2,000	2,900	4,200
<b>Austria</b>	11,000	15,000	21,000	5,800	7,900	11,000
<b>Azerbaijan</b>	460	680	960	1,600	2,200	3,200
<b>Bahamas, The</b>	86	130	170	40	60	80
<b>Bahrain</b>	330	510	750	200	300	400
<b>Bangladesh</b>	9,100	14,000	18,000	67,000	96,000	130,000
<b>Barbados</b>	12	18	25	10	20	30
<b>Belarus</b>	2,400	3,200	4,400	7,800	10,000	14,000
<b>Belgium</b>	8,500	12,000	18,000	5,000	6,800	9,500
<b>Belize</b>	7	11	15	20	30	40
<b>Benin</b>	24	39	55	220	350	520
<b>Bermuda</b>	29	44	59	9	10	20
<b>Bhutan</b>	31	54	79	150	240	350
<b>Bolivia</b>	28	59	100	130	250	450
<b>Bosnia &amp; Herzegovina</b>	680	950	1,200	2,300	3,100	3,900
<b>Botswana</b>	57	91	140	100	150	200
<b>Brazil</b>	9,800	14,000	22,000	20,000	28,000	43,000
<b>Brunei Darussalam</b>	35	53	76	20	30	40
<b>Bulgaria</b>	2,900	3,900	5,200	6,800	9,000	12,000
<b>Burkina Faso</b>	59	95	130	600	900	1,300
<b>Burundi</b>	7	11	17	200	320	500
<b>Cabo Verde</b>	6	9	13	30	40	50
<b>Cambodia</b>	240	360	520	1,900	2,800	4,100

Impact attributed to fossil fuel-related air pollution by country/region globally						
Country/Region	Estimated total cost (Million USD)			Estimated total premature deaths (2018)		
	Low	Central	High	Low	Central	High
<b>Cameroon</b>	99	160	230	570	900	1,000
<b>Canada</b>	25,000	38,000	57,000	15,000	21,000	30,000
<b>Central African Republic</b>	11	18	27	160	260	400
<b>Chad</b>	61	99	140	550	850	1,000
<b>Chile</b>	1,600	2,600	4,500	2,300	3,800	6,600
<b>China Mainland</b>	650,000	900,000	1,200,000	1,300,000	1,800,000	2,500,000
<b>Colombia</b>	1,500	2,400	3,600	4,800	6,900	9,800
<b>Comoros</b>	1	2	3	9	10	19
<b>Congo, Dem. Rep.</b>	79	130	210	1,100	2,000	3,300
<b>Congo, Rep.</b>	9	15	22	44	70	100
<b>Costa Rica</b>	230	340	450	380	530	710
<b>Côte d'Ivoire</b>	16	38	66	90	200	400
<b>Croatia</b>	2,000	2,800	3,700	3,300	4,400	5,700
<b>Cuba</b>	No Data	No Data	No Data	2,000	2,800	3,800
<b>Cyprus</b>	380	570	790	310	440	630
<b>Czech Republic</b>	7,700	11,000	15,000	8,000	11,000	14,000
<b>Denmark</b>	4,600	6,700	9,500	2,000	2,800	3,800
<b>Djibouti</b>	No Data	No Data	No Data	40	80	100
<b>Dominica</b>	2	3	4	5	7	9
<b>Dominican Republic</b>	310	490	700	600	1,000	1,400
<b>Ecuador</b>	670	1,000	1,500	2,000	2,700	3,700
<b>Egypt, Arab Rep.</b>	4,400	6,900	10,000	22,000	32,000	51,000
<b>El Salvador</b>	160	260	370	697	1,100	1,500
<b>Equatorial Guinea</b>	3	7	12	4	9	20
<b>Eritrea</b>	39	66	98	390	640	970
<b>Estonia</b>	430	650	900	460	630	820
<b>Eswatini</b>	41	66	97	120	180	270
<b>Ethiopia</b>	240	370	480	2,600	3,900	5,400
<b>Fiji</b>	8	12	17	20	20	30
<b>Finland</b>	2,800	4,300	5,800	1,600	2,100	2,700
<b>France</b>	37,000	54,000	79,000	27,000	37,000	55,000
<b>Gabon</b>	5	9	13	8	10	20



Impact attributed to fossil fuel-related air pollution by country/region globally						
Country/Region	Estimated total cost (Million USD)			Estimated total premature deaths (2018)		
	Low	Central	High	Low	Central	High
<b>Gambia, The</b>	4	8	11	60	100	150
<b>Georgia</b>	510	700	890	2,200	2,900	3,600
<b>Germany</b>	94,000	140,000	210,000	57,000	81,000	120,000
<b>Ghana</b>	38	73	120	200	370	600
<b>Greece</b>	4,200	6,100	8,700	6,000	8,400	12,000
<b>Greenland</b>	No Data	No Data	No Data	8	10	20
<b>Grenada</b>	3	4	6	6	8	10
<b>Guam</b>	No Data	No Data	No Data	10	16	22
<b>Guatemala</b>	390	620	870	1,200	1,700	2,400
<b>Guinea</b>	27	43	60	270	410	590
<b>Guinea Bissau</b>	3	6	8	40	60	90
<b>Guyana</b>	6	9	12	20	20	30
<b>Haiti</b>	61	100	150	772	1,300	1,900
<b>Honduras</b>	100	170	260	600	980	2,000
<b>Hungary</b>	6,700	9,400	13,000	9,400	13,000	17,000
<b>Iceland</b>	76	110	150	30	40	50
<b>India</b>	100,000	150,000	190,000	715,000	1,000,000	1,300,000
<b>Indonesia</b>	7,600	11,000	16,000	30,000	44,000	61,000
<b>Iran, Islamic Rep.</b>	3,800	5,300	7,300	13,000	17,000	24,000
<b>Iraq</b>	1,400	2,100	2,800	2,500	3,500	4,800
<b>Ireland</b>	2,500	3,800	5,000	900	1,200	1,600
<b>Israel</b>	3,000	4,500	6,200	1,800	2,500	3,600
<b>Italy</b>	41,000	61,000	91,000	39,000	56,000	83,000
<b>Jamaica</b>	90	140	200	310	460	640
<b>Japan</b>	88,000	130,000	180,000	75,000	100,000	150,000
<b>Jordan</b>	300	490	700	800	1,200	1,900
<b>Kazakhstan</b>	2,000	2,800	3,900	3,500	4,800	6,400
<b>Kenya</b>	190	290	380	1,100	1,600	2,300
<b>Kiribati</b>	0	0	0	0	0	0
<b>Korea, Dem. People's Rep.</b>	No Data	No Data	No Data	24,000	38,000	56,000
<b>Korea, Rep.</b>	37,000	56,000	85,000	28,000	40,000	61,000
<b>Kosovo</b>	180	270	380	800	1,200	1,700

Impact attributed to fossil fuel-related air pollution by country/region globally						
Country/Region	Estimated total cost (Million USD)			Estimated total premature deaths (2018)		
	Low	Central	High	Low	Central	High
<b>Kuwait</b>	840	1,300	1,700	290	410	600
<b>Kyrgyz Republic</b>	77	110	150	800	1,100	1,600
<b>Lao PDR</b>	320	510	720	1,400	2,000	2,900
<b>Latvia</b>	850	1,200	1,700	1,100	1,500	2,100
<b>Lebanon</b>	890	1,400	2,100	1,800	2,700	4,200
<b>Lesotho</b>	39	67	100	700	588	870
<b>Liberia</b>	2	3	4	20	40	60
<b>Libya</b>	300	470	660	600	900	1,300
<b>Lithuania</b>	1,700	2,300	3,000	2,000	2,600	3,300
<b>Luxembourg</b>	1,000	1,500	2,300	250	350	500
<b>Madagascar</b>	11	19	29	180	300	450
<b>Malawi</b>	6	11	16	170	280	410
<b>Malaysia</b>	2,800	4,500	6,700	4,300	6,600	10,000
<b>Maldives</b>	17	26	38	30	40	60
<b>Mali</b>	58	99	140	500	800	1,300
<b>Malta</b>	140	200	260	120	170	220
<b>Marshall Islands</b>	0	1	1	1	2	3
<b>Mauritania</b>	11	20	28	100	170	250
<b>Mauritius</b>	23	33	44	30	50	60
<b>Mexico</b>	20,000	29,000	41,000	37,000	51,000	73,000
<b>Micronesia, Fed. Sts.</b>	1	2	3	4	6	9
<b>Moldova</b>	450	600	750	2,300	3,000	3,600
<b>Mongolia</b>	130	200	270	410	570	780
<b>Montenegro</b>	150	210	280	370	480	620
<b>Morocco</b>	670	1,100	1,600	3,300	5,100	7,500
<b>Mozambique</b>	12	20	30	200	340	530
<b>Myanmar</b>	1,300	1,900	2,500	12,000	18,000	24,000
<b>Namibia</b>	15	26	39	36	56	84
<b>Nepal</b>	580	940	1,400	7,800	12,000	18,000
<b>Netherlands</b>	14,000	21,000	30,000	7,200	9,900	14,000
<b>New Zealand</b>	190	270	350	110	140	170
<b>Nicaragua</b>	38	60	83	300	440	600

Impact attributed to fossil fuel-related air pollution by country/region globally						
Country/Region	Estimated total cost (Million USD)			Estimated total premature deaths (2018)		
	Low	Central	High	Low	Central	High
<b>Niger</b>	35	60	89	530	880	1,000
<b>Nigeria</b>	1,300	2,200	3,200	4,600	7,600	13,000
<b>North Macedonia</b>	390	540	700	1,200	1,600	2,000
<b>Northern Mariana Islands</b>	No Data	No Data	No Data	3	4	6
<b>Norway</b>	4,300	6,000	8,500	1,500	2,100	2,800
<b>Oman</b>	200	320	430	140	210	300
<b>Pakistan</b>	3,800	6,100	9,200	32,000	50,000	76,000
<b>Palestine</b>	82	120	160	400	500	700
<b>Panama</b>	170	260	360	230	310	410
<b>Papua New Guinea</b>	75	120	200	290	430	620
<b>Paraguay</b>	85	140	210	250	380	540
<b>Peru</b>	550	970	1,700	1,500	2,500	4,600
<b>Philippines</b>	2,500	4,000	6,000	11,000	17,000	27,000
<b>Poland</b>	21,000	29,000	38,000	30,000	39,000	51,000
<b>Portugal</b>	2,700	4,100	6,300	3,300	4,800	7,200
<b>Puerto Rico</b>	470	730	1,000	420	610	860
<b>Qatar</b>	1,000	1,600	2,400	140	230	410
<b>Romania</b>	9,100	13,000	17,000	17,000	22,000	29,000
<b>Russian Federation</b>	50,000	68,000	97,000	89,000	120,000	160,000
<b>Rwanda</b>	18	30	45	240	390	590
<b>Samoa</b>	0	0	1	1	1	2
<b>São Tomé and Príncipe</b>	1	1	2	4	7	10
<b>Saudi Arabia</b>	3,800	6,000	8,800	2,200	3,300	5,000
<b>Senegal</b>	70	120	160	480	743	1,100
<b>Serbia</b>	2,600	3,700	4,900	8,500	11,000	15,000
<b>Seychelles</b>	6	8	11	6	9	11
<b>Sierra Leone</b>	8	13	18	120	190	280
<b>Singapore</b>	2,500	4,000	6,500	890	1,000	2,000
<b>Slovak Republic</b>	3,700	5,100	6,700	4,100	5,400	7,000
<b>Slovenia</b>	1,300	1,800	2,500	1,300	1,700	2,300
<b>Solomon Islands</b>	2	4	6	10	20	20
<b>Somalia</b>	No Data	No Data	No Data	610	1,000	2,000

Impact attributed to fossil fuel-related air pollution by country/region globally						
Country/Region	Estimated total cost (Million USD)			Estimated total premature deaths (2018)		
	Low	Central	High	Low	Central	High
<b>South Africa</b>	4,300	6,300	8,200	9,700	13,000	16,000
<b>South Sudan</b>	53	87	130	290	480	750
<b>Spain</b>	16,000	24,000	36,000	17,000	25,000	37,000
<b>Sri Lanka</b>	460	760	1,100	2,100	3,300	4,800
<b>St. Lucia</b>	5	8	11	10	10	20
<b>St. Vincent &amp; the Grenadines</b>	2	3	4	5	7	10
<b>Sudan</b>	180	320	460	1,900	3,100	4,600
<b>Suriname</b>	5	8	11	10	20	30
<b>Sweden</b>	5,400	7,800	11,000	3,000	4,000	5,200
<b>Switzerland</b>	11,000	16,000	23,000	4,100	5,500	7,600
<b>Syrian Arab Republic</b>	No Data	No Data	No Data	3,100	4,700	7,100
<b>Taiwan</b>	11,000	16,000	23,000	12,000	16,000	24,000
<b>Tajikistan</b>	65	96	130	770	1,000	2,000
<b>Tanzania</b>	76	130	180	700	1,000	1,500
<b>Thailand</b>	7,000	11,000	15,000	17,000	24,000	34,000
<b>TimorLeste</b>	1	3	5	10	20	30
<b>Togo</b>	11	18	26	100	200	300
<b>Tonga</b>	0	1	1	1	1	2
<b>Trinidad and Tobago</b>	41	67	100	40	70	100
<b>Tunisia</b>	240	400	590	1,300	2,100	3,100
<b>Turkey</b>	14,000	21,000	30,000	28,000	40,000	58,000
<b>Turkmenistan</b>	240	360	520	500	700	1,000
<b>Uganda</b>	36	57	79	500	700	1,100
<b>Ukraine</b>	6,000	8,000	10,000	35,000	45,000	57,000
<b>United Arab Emirates</b>	3,500	5,900	9,400	900	1,500	2,400
<b>United Kingdom</b>	46,000	66,000	98,000	30,000	41,000	62,000
<b>United States</b>	430,000	610,000	870,000	170,000	230,000	310,000
<b>Uruguay</b>	280	450	730	800	630	1,000
<b>Uzbekistan</b>	410	590	810	3,400	4,800	6,600
<b>Vanuatu</b>	1	2	3	5	7	11
<b>Venezuela, RB</b>	1,800	2,800	4,200	1,900	2,900	4,200
<b>Vietnam</b>	4,500	6,800	9,800	28,000	41,000	58,000

Impact attributed to fossil fuel-related air pollution by country/region globally						
Country/Region	Estimated total cost (Million USD)			Estimated total premature deaths (2018)		
	Low	Central	High	Low	Central	High
Virgin Islands (U.S.)	No Data	No Data	No Data	10	20	20
Palestine	80	120	160	400	500	700
Yemen, Rep.	150	280	450	1,800	3,100	5,200
Zambia	33	56	80	200	300	500
Zimbabwe	56	91	130	200	300	500

## ENDNOTES

- Strasert, B., Teh, S. C. & Cohan, D. S. Air quality and health benefits from potential coal power plant closures in Texas. *J. Air & Waste Manage.* 69, 333–350 (2019).
- Lelieveld, J., et al. Effects of fossil fuel and total anthropogenic emission removal on public health and climate. *PNAS* 116, 7192–7197 (2019).
- Watts N., et al. Health and climate change: Policy responses to protect public health. *Lancet* 386, 1861–1914. (2015).
- Lelieveld, J., et al. Effects of fossil fuel and total anthropogenic emission removal on public health and climate. *PNAS* 116, 7192–7197 (2019).
- Watts N., et al. Health and climate change: Policy responses to protect public health. *Lancet* 386, 1861–1914 (2015).
- Anderson, H., et al. Quantitative systematic review of short term associations between ambient air pollution (particulate matter, ozone, nitrogen dioxide, sulphur dioxide and carbon monoxide), and mortality and morbidity. London, Department of Health. Available at: <https://www.gov.uk/government/publications/quantitative-systematic-review-of-short-term-associations-between-ambient-air-pollution-particulate-matter-ozone-nitrogen-dioxide-sulphur-dioxide-and-carbon-monoxide-and-mortality-and-morbidity> [Accessed January 9, 2020].
- Guarnieri, M. & Balmes, J. Outdoor air pollution and asthma. *Lancet* 383, 1581–1592 (2014).
- United States Environmental Protection Agency. Ground-level Ozone Pollution. Available at: <https://www.epa.gov/ground-level-ozone-pollution/ground-level-ozone-basics> (2020) [Accessed January 9, 2020].
- World Health Organization. Review of evidence on health aspects of air pollution – REVIHAAP Project Technical Report. WHO: Bonn (2013). Available at: [http://www.euro.who.int/\\_data/assets/pdf\\_file/0004/193108/REVIHAAP-Final-technical-report-final-version.pdf?ua=1](http://www.euro.who.int/_data/assets/pdf_file/0004/193108/REVIHAAP-Final-technical-report-final-version.pdf?ua=1) [Accessed January 16, 2020].
- Morakinyo, O., Mokgobu, M., Mukhola, M. & Hunter, R. Health outcomes of exposure to biological and chemical components of inhalable and respirable particulate matter. *Int. J. Environ. Res. Public Health* 13, 592 (2016). DOI: 10.3390/ijerph13060592.
- Cassee, F.R., Héroux, M.-E., Gerlofs-Nijland, M.E. & Kelly, F.J. Particulate matter beyond mass: recent health evidence on the role of fractions, chemical constituents and sources of emission. *Inhal. Toxicol.* 25, 802–812 (2013). DOI: 10.3109/08958378.2013.850127.
- Morakinyo, O., Mokgobu, M., Mukhola, M. & Hunter, R. Health outcomes of exposure to biological and chemical components of inhalable and respirable particulate matter. *Int. J. Environ. Res. Public Health* 13, 592 (2016). DOI: 10.3390/ijerph13060592.
- WHO, 2006. World Health Organisation. Air Quality Guidelines Global Update 2005. Available at: [http://www.euro.who.int/\\_data/assets/pdf\\_file/0005/78638/E90038.pdf?ua=1](http://www.euro.who.int/_data/assets/pdf_file/0005/78638/E90038.pdf?ua=1) [Accessed January 16, 2020].
- Lelieveld, J., Evans, J., Fnais, M., Giannadaki, D. & Pozzer, A. The contribution of outdoor air pollution sources to premature mortality on a global scale. *Nature* 25, 367–371 (2015). DOI: 10.1038/nature15371
- Zyrichidou, I., Koukouli, M., Balis, D., Markakis, K., Poupkou, A., Katragkou, E., Kioutsioukis, I., Melas, D., Boersma, K., van Roozendaal, M. Identification of surface NOx emission sources on a regional scale using OMI NO2. *Atmos. Environ.* 101, 82–93 (2015). DOI: 10.1016/j.atmosenv.2014.11.023
- Squizzato, S., Cazzaro, M., Innocente, E., Visin, F., Hopke, P. & Rampazzo, G. Urban air quality in a mid-size city — PM2.5 composition, sources and identification of impact areas: From local to long range contributions. *Atmos. Res.* 186, 51–62 (2017). DOI: 10.1016/J.ATMOSRES.2016.11.011 <https://www.epa.gov/no2-pollution/basic-information-about-no2#What%20is%20NO2>
- IRENA. Renewable Power Generation Costs in 2018. International Renewable Energy Agency, Abu Dhabi (2019).
- Farfan, J. and Breyer, C. Structural changes of global power generation capacity towards sustainability and the risk of stranded investments supported by a sustainability indicator. *Journal of Cleaner Production* 14, 370–384 (2017).
- Wang, B., Xu, D., Jing, Z., Liu, D., Yan, S. & Wang, Y. Effect of long-term exposure to air pollution on type 2 diabetes mellitus risk: a systemic review and meta-analysis of cohort studies. *Eur. J. Endocrinol.* 171, R173–R182 (2014).
- Cohen, A. J. et al. Estimates and 25-year trends of the global burden of disease attributable to ambient air pollution: an analysis of data from the Global Burden of Diseases Study 2015. *Lancet* 389, 1907–1918 (2017).
- Han, M.-H., et al. Association between hemorrhagic stroke occurrence and meteorological factors and pollutants. *BMC Neurol.* 16, 59 (2016).
- Sunyer, J. & Davvand, P. Pre-natal brain development as a target for urban air pollution. *Basic Clin. Pharmacol. Toxicol.* 125, Suppl 3, 81–88 (2019).
- Lelieveld, J., Evans, J. S., Fnai, M., Giannadaki, D. & Pozzer, A. The contribution of outdoor air pollution sources to premature mortality on a global scale. *Nature* 525, 367–371 (2015).
- Park, M. et al. Differential toxicities of fine particulate matters from various sources. *Sci. Rep.* 8, 17007 (2018).
- Li, J. et al. Differing toxicity of ambient particulate matter (PM) in global cities. *Atmos. Environ.* 212, 305–315 (2019).
- Lelieveld, J., Evans, J. S., Fnai, M., Giannadaki, D. & Pozzer, A. The contribution of outdoor air pollution sources to premature mortality on a global scale. *Nature* 525, 367–371 (2015).
- Lelieveld, J., Evans, J. S., Fnai, M., Giannadaki, D. & Pozzer, A. The contribution of outdoor air pollution sources to premature mortality on a global scale. *Nature* 525, 367–371 (2015).
- World Health Organization, 2006. WHO Air quality guidelines for particulate matter, ozone, nitrogen dioxide and sulfur dioxide: global update 2005: summary of risk assessment (No. WHO/SDE/PHE/OEH/06.02). Geneva: World Health Organization.
- World Health Organization (WHO). “WHO expert consultation: available evidence for the future update of the WHO global air quality guidelines (AQGs).” WHO: Geneva, Switzerland (2016).
- WHO. ‘Air pollution: Ambient air pollution - a major threat to health and climate.’ World Health Organization, 2020. Available at: <https://www.who.int/airpollution/ambient/en/> [Accessed January 8 2020].
- Shindell, D. & Smith, C.J. Climate and air-quality benefits of a realistic phase-out of fossil fuels. *Nature* 573, 408–411 (2019).
- Watts N., et al. Health and climate change: Policy responses to protect public health. *Lancet* 386, 1861–1914 (2015).
- Williams, M.L., et al. Public health air pollution impacts of pathway options to meet the 2050 UK Climate Change Act target: a modelling study. *Public Health Research* 6.7 (2018).
- Lelieveld, J., et al. Effects of fossil fuel and total anthropogenic emission removal on public health and climate. *PNAS* 116, 7192–7197 (2019).
- Larkin, A., et al. Global land use regression model for nitrogen dioxide air pollution. *Environmental science & technology* 51.12: 6957–6964. (2017)
- World Health Organization. Health risks of air pollution in Europe—HRAPIE project recommendations for concentration–response functions for cost–benefit analysis of particulate matter, ozone and nitrogen dioxide. UN City: Copenhagen, Denmark (2013).
- Burnett, R., et al. Global estimates of mortality associated with long-term exposure to outdoor fine particulate matter. *Proceedings of the National Academy of Sciences* 115.38: 9592–9597. (2018).
- Malley CS, Daven K. Henze, Johan C.I. Kuylenstierna, Harry W. Vallack, Yanko Davila, Susan C. Anenberg, Michelle C. Turner, and Mike R. Ashmore 2017: Updated Global Estimates of Respiratory Mortality in Adults ≥30 Years of Age Attributable to Long-Term Ozone Exposure. *Environmental Health Perspectives* 125:8 CID: 087021 <https://doi.org/10.1289/EHP1390>
- Silver, B., et al. “Substantial changes in air pollution across China during 2015–2017.” *Environmental Research Letters* 13.11 (2018): 114012.
- Shindell, D. & Smith, C.J. Climate and air-quality benefits of a realistic phase-out of fossil fuels. *Nature* 573, 408–411 (2019).
- Watts, N. et al. Health and climate change: policy responses to protect public health. *The Lancet* 386, 1861–1914 (2015).
- United States Environmental Protection Agency: Office of Air and Radiation. The benefits and costs of the Clean Air Act from 1990 to 2020. Available at: [https://www.epa.gov/sites/production/files/2015-07/documents/fullreport\\_rev\\_a.pdf](https://www.epa.gov/sites/production/files/2015-07/documents/fullreport_rev_a.pdf) (2011) [Accessed January 9, 2020].
- Landrigan, P. J. et al. The Lancet Commission on pollution and health. *The Lancet* 391, 10119, 462–512 (2018).
- <https://www.greenpeace.org/international/story/24312/the-future-of-transport-is-zero-carbon/>
- Teske, S. et al. the Energy [R]evolution (5th edition). Greenpeace International, Global Wind Energy Council & Solar Power Europe (2015). Available at: [https://www.duesseldorf.greenpeace.de/sites/www.duesseldorf.greenpeace.de/files/greenpeace\\_energy-revolution\\_erneuerbare\\_2050\\_20150921.pdf](https://www.duesseldorf.greenpeace.de/sites/www.duesseldorf.greenpeace.de/files/greenpeace_energy-revolution_erneuerbare_2050_20150921.pdf) [Accessed January 8, 2020].
- Watts N., et al. Health and climate change: Policy responses to protect public health. *Lancet* 386, 1861–1914. (2015).

- 49 Keegan, M. 'Shenzhen's silent revolution: world's first fully electric bus fleet quietens Chinese megacity'. The Guardian December 12, 2018. Available at: <https://www.london.gov.uk/what-we-do/environment/pollution-and-air-quality/cleaner-buses> [Accessed January 8, 2020].
- 50 Keegan, M. 'Shenzhen's silent revolution: world's first fully electric bus fleet quietens Chinese megacity'. The Guardian. December 12, 2018. Available at: <https://www.london.gov.uk/what-we-do/environment/pollution-and-air-quality/cleaner-buses> [Accessed January 8, 2020].
- 51 [https://new.mta.info/system\\_modernization/sustainabletransit](https://new.mta.info/system_modernization/sustainabletransit)
- 52 <https://www.oslo.kommune.no/politics-and-administration/green-oslo/best-practices/car-free-city/#gref>
- 53 Nieuwenhuijsen, M. J. & Khreis, H. Car free cities: Pathway to healthy urban living. *Environ. Int.* 94, 251–262 (2016).
- 54 Masiol, M. Thirteen years of air pollution hourly monitoring in a large city: Potential sources, trends, cycles and effects of car-free days. *Sci. Total Environ.* 494–495, 84–96 (2014).
- 55 [https://www.greenpeace.de/sites/www.greenpeace.de/files/publications/gp-cleanairnow\\_carindustryreport\\_full\\_v5\\_0919\\_72ppi\\_0.pdf](https://www.greenpeace.de/sites/www.greenpeace.de/files/publications/gp-cleanairnow_carindustryreport_full_v5_0919_72ppi_0.pdf)
- 56 [https://storage.googleapis.com/planet4-international-stateless/2018/01/1b96c158-air\\_pollution-transport\\_report-2018.pdf](https://storage.googleapis.com/planet4-international-stateless/2018/01/1b96c158-air_pollution-transport_report-2018.pdf)
- 57 Montes, F. et al. Do Health Benefits Outweigh the Costs of Mass Recreational Programs? An Economic Analysis of Four Ciclovía Programs. *J. Urban Health* 89, 153–170 (2012).
- 58 Triana, C. A. et al. Active streets for children: The case of the Bogotá Ciclovía. *PLoS One* 14, e0207791 (2019).
- 59 Montes, F. et al. Do Health Benefits Outweigh the Costs of Mass Recreational Programs? An Economic Analysis of Four Ciclovía Programs. *J. Urban Health* 89, 153–170 (2012).
- 60 Engelberg, J. A., et al. Ciclovía Participation and Impacts in San Diego, CA: The First CicloSDias. *Prev Med.* 69, S66–S73 (2014).
- 61 Lelieveld, J., et al. Effects of fossil fuel and total anthropogenic emission removal on public health and climate. *PNAS* 116, 7192–7197 (2019).
- 62 Millstein, D., Wisner, R., Bolinger, M. & Barbose, G. The climate and air-quality benefits of wind and solar power in the United States. *Nature Energy* 2, 17134 (2017).
- 63 Strasert, B., Teh, S. C. & Cohan, D. S. Air quality and health benefits from potential coal power plant closures in Texas. *J. Air & Waste Manage.* 69, 333–350 (2019).
- 64 Millstein, D., Wisner, R., Bolinger, M. & Barbose, G. The climate and air-quality benefits of wind and solar power in the United States. *Nature Energy* 2, 17134 (2017).
- 65 Kalia, V., Perera, F. & Tang, D. Environmental Pollutants and Neurodevelopment: Review of Benefits From Closure of a Coal-Burning Power Plant in Tongliang, China. *Global Pediatric Health* (2017).
- 66 Russell, M. C., Belle, J. H. & Liu, Y. The impact of three recent coal-fired power plant closings on Pittsburgh air quality: A natural experiment. *Journal of the Air & Waste Management Association*, 67, 3–16 (2017).
- 67 Dockery, D.W., et al. Effect of air pollution control on mortality and hospital admissions in Ireland. Research Report 176. Health Effects Institute, Boston, MA (2013).
- 68 Strasert, B., Teh, S. C. & Cohan, D. S. Air quality and health benefits from potential coal power plant closures in Texas. *J. Air & Waste Manage.* 69, 333–350 (2019).
- 69 Lelieveld, J., Evans, J., Fnais, M. et al. The contribution of outdoor air pollution sources to premature mortality on a global scale. *Nature* 525, 367–371 (2015).
- 70 Larkin, A., et al. Global land use regression model for nitrogen dioxide air pollution. *Environmental science & technology* 51.12: 6957–6964. (2017)
- 71 Levy, R., et al. MODIS Atmosphere L2 Aerosol Product. NASA MODIS Adaptive Processing System, Goddard Space Flight Center, USA: [http://dx.doi.org/10.5067/MODIS/MOD04\\_L2.006](http://dx.doi.org/10.5067/MODIS/MOD04_L2.006) (2015). f
- 72 GBD 2017 Mortality Collaborators. Global, regional, and national age-sex-specific mortality and life expectancy, 1950–2017: a systematic analysis for the Global Burden of Disease Study 2017. *The Lancet.* 392:1684–735. (2018).
- 73 Anenberg, Susan C., et al. Estimates of the Global Burden of Ambient PM2.5, Ozone, and NO2 on Asthma Incidence and Emergency Room Visits. *Environmental health perspectives* 126.10: 107004. (2018).
- 74 Chawanpaiboon S, Vogel JP, Moller AB, Lumbiganon P, Petzold M, Hogan D, Landoulsi S, Jampathong N, Kongwattanakul K, Laopaiboon M, Lewis C, Rattanakonchai S, Teng DN, Thinkhamrop J, Watananirun K, Zhang J, Zhou W, Gülmezoglu AM 2019: Global, regional, and national estimates of levels of preterm birth in 2014: a systematic review and modelling analysis. *Lancet Glob Health* 7(1):e37–e46. doi: 10.1016/S2214-109X(18)30451-0
- 75 World Bank: <https://datahelpdesk.worldbank.org/knowledgebase/articles/906519-world-bank-country-and-lending-groups>
- 76 Burnett, R, et al. Global estimates of mortality associated with long-term exposure to outdoor fine particulate matter. *Proceedings of the National Academy of Sciences* 115.38: 9592–9597. (2018).
- 77 Burnett R et al 2018: Global estimates of mortality associated with long-term exposure to outdoor fine particulate matter. *Proceedings of the National Academy of Sciences* Sep 2018, 115 (38) 9592–9597; DOI: 10.1073/pnas.1803222115
- 78 Cohen, A. J. et al. Estimates and 25-year trends of the global burden of disease attributable to ambient air pollution: an analysis of data from the Global Burden of Diseases Study 2015. *Lancet* 389, 1907–1918 (2017).
- 79 European Environment Agency (EEA) 2014: Costs of air pollution from European industrial facilities 2008–2012 — an updated assessment. EEA Technical report No 20/2014. <https://www.eea.europa.eu/publications/costs-of-air-pollution-2008-2012>
- 80 OECD, The Economic Consequences of Outdoor Air Pollution, OECD Publishing, Paris. <http://dx.doi.org/10.1787/9789264257474-en> (2016)
- 81 Cohen, A. J. et al. Estimates and 25-year trends of the global burden of disease attributable to ambient air pollution: an analysis of data from the Global Burden of Diseases Study 2015. *Lancet* 389, 1907–1918 (2017).
- 82 Birchby D, Stedman J, Whiting S, Vedrenne M: Air Quality damage cost update 2019. Report for Defra. AQ0650. Ricardo Energy & Environment, United Kingdom. (2019)
- 83 Brandt SJ, Perez L, Künzli N, Lurmann F, McConnell R 2012: Costs of childhood asthma due to traffic-related pollution in two California communities. *European Respiratory Journal* Aug 2012, 40 (2) 363–370; DOI: 10.1183/09031936.00157811.
- 84 Anenberg, Susan C., et al. Estimates of the Global Burden of Ambient PM2.5, Ozone, and NO2 on Asthma Incidence and Emergency Room Visits. *Environmental health perspectives* 126.10: 107004. (2018).
- 85 Trasande L, Malecha P, Attina TM 2016: Particulate Matter Exposure and Preterm Birth: Estimates of U.S. Attributable Burden and Economic Costs. *Environmental Health Perspectives* 124:12. <https://doi.org/10.1289/ehp.1510810>
- 86 World Health Organization. Health risks of air pollution in Europe—HRAPIE project recommendations for concentration–response functions for cost–benefit analysis of particulate matter, ozone and nitrogen dioxide. UN City: Copenhagen, Denmark (2013).
- 87 European Environment Agency (EEA) 2014: Costs of air pollution from European industrial facilities 2008–2012 — an updated assessment. EEA Technical report No 20/2014. <https://www.eea.europa.eu/publications/costs-of-air-pollution-2008-2012>
- 88 Zheng, Xue-yan, et al. “Association between air pollutants and asthma emergency room visits and hospital admissions in time series studies: a systematic review and meta-analysis.” *PloS one* 10.9 (2015).
- 89 Trasande L, Malecha P, Attina TM 2016: Particulate Matter Exposure and Preterm Birth: Estimates of U.S. Attributable Burden and Economic Costs. *Environmental Health Perspectives* 124:12. <https://doi.org/10.1289/ehp.1510810>
- 90 Davdand P et al. 2013: Maternal Exposure to Particulate Air Pollution and Term Birth Weight: A Multi-Country Evaluation of Effect and Heterogeneity. *Environmental Health Perspectives.* [https://ehp.niehs.nih.gov/doi/full/10.1289/ehp.1205575?url\\_ver=Z39.88-2003&rfr\\_id=ori:rid:crossref.org&rfr\\_dat=cr\\_pub%3dpubmed](https://ehp.niehs.nih.gov/doi/full/10.1289/ehp.1205575?url_ver=Z39.88-2003&rfr_id=ori:rid:crossref.org&rfr_dat=cr_pub%3dpubmed)
- 91 Malley CS,Daven K. Henze,Johan C.I. Kuylenstierna,Harry W. Vallack,Yanko Davila,Susan C. Anenberg,Michelle C. Turner,and Mike R. Ashmore 2017: Updated Global Estimates of Respiratory Mortality in Adults ≥30Years of Age Attributable to Long-Term Ozone Exposure. *Environmental Health Perspectives* 125:8 CID: 087021 <https://doi.org/10.1289/EHP1390>
- 92 World Health Organization. Health risks of air pollution in Europe—HRAPIE project recommendations for concentration–response functions for cost–benefit analysis of particulate matter, ozone and nitrogen dioxide. UN City: Copenhagen, Denmark (2013).
- 93 Achakulwisut, Pattanun, et al. “Global, national, and urban burdens of paediatric asthma incidence attributable to ambient NO2 pollution: estimates

from global datasets” *The Lancet Planetary Health* 3.4 (2019): e166–e178.

94 Anenberg, Susan C., et al. Estimates of the Global Burden of Ambient PM2.5, Ozone, and NO2 on Asthma Incidence and Emergency Room Visits. *Environmental health perspectives* 126.10: 107004. (2018).

95 Chawanpaiboon S, Vogel JP, Moller AB, Lumbiganon P, Petzold M, Hogan D, Landoulsi S, Jampathong N, Kongwattanakul K, Laopaiboon M, Lewis C, Rattanakonchai S, Teng DN, Thinkhamrop J, Watananirun K, Zhang J, Zhou W, Gülmezoglu AM 2019: Global, regional, and national estimates of levels of preterm birth in 2014: a systematic review and modelling analysis. *Lancet Glob Health* 7(1):e37–e46. doi: 10.1016/S2214-109X(18)30451-0

96 World Bank : <https://datahelpdesk.worldbank.org/knowledgebase/articles/906519-world-bank-country-and-lending-groups>

97 GBD 2017 Mortality Collaborators. Global, regional, and national age-sex-specific mortality and life expectancy, 1950–2017: a systematic analysis for the Global Burden of Disease Study 2017. *The Lancet.* 392:1684–735. (2018).

98 European Environment Agency (EEA) 2014: Costs of air pollution from European industrial facilities 2008–2012 — an updated assessment. EEA Technical report No 20/2014. <https://www.eea.europa.eu/publications/costs-of-air-pollution-2008-2012>

Greenpeace is an independent global campaigning organisation that acts to change attitudes and behaviour, to protect and conserve the environment and to promote peace.



**Greenpeace MENA**

Berytech Building,  
Technology Pole.  
Damascus Road, Mathaf, Beirut, Lebanon  
k.v.k. reg 72184981  
info.arabic@greenpeace.org  
www.greenpeace.org/mena